(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 29 January 2004 (29.01.2004)

PCT

(10) International Publication Number WO 2004/009823 A1

(51) International Patent Classification7: 15/90

C12N 15/85,

(21) International Application Number:

PCT/EP2003/007946

(22) International Filing Date:

21 July 2003 (21.07.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0216648.6

GB 19 July 2002 (19.07.2002)

(71) Applicant (for all designated States except US): LONZA BIOLOGICS PLC. [GB/GB]; 228 Bath Road, Slough SL1 4DY, Berkshire (GB).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): KALLMEIER, Robert [GB/GB]; 47 Fox Road, Holmer Green, Bucks HP 156SF (GB). GAY, Robert [GB/GB]; 7 Durlgy Avenue, Pinner Middx HA 51JQ (GB).
- (74) Common Representative: LONZA BIOLOGICS PLC.; Legal Department, 228 Bath Road, Slough SL1 4DY, Berkshire (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, $\mathsf{MX}, \mathsf{MZ}, \mathsf{NI}, \mathsf{NO}, \mathsf{NZ}, \mathsf{OM}, \mathsf{PG}, \mathsf{PH}, \mathsf{PL}, \mathsf{PT}, \mathsf{RO}, \mathsf{RU}, \mathsf{SC},$

SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG. KP. KR. KZ. LC. LK. LR. LS. LT. LU. LV. MA. MD. MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH. GM. KE, LS, MW. MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ. TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK. EE. ES. FI. FR. GB. GR, HU. IE. IT. LU, MC. NL, PT. RO. SE. SI. SK. TR), OAPI patent (BF, BJ. CF, CG, CI, CM, GA, GN, GQ, GW, ML. MR. NE. SN, TD. TG)
- of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD OF EXPRESSING RECOMBINANT PROTEIN IN CHO CELLS

(57) Abstract: Method of expressing recombinant protein in CHO cells, by using an expression vector comprising the murine IgG 2A gene locus.

10

15

20

25

30



Method of expressing recombinant protein in CHO cells

The present invention relates to a method for expressing a recombinant product gene in a CHO cell line as well as to recombinant CHO host cells and to novel expression vector constructs.

The Chinese Hamster ovary cell (CHO) mammalian expression system is widely used in production of recombinant protein. Apart from lymphoid cell lines such as hybridoma cell lines, it is one of the few cell types allowing for simple and efficient high-density suspension batch culture of animal cell. Furthermore, they allow for very high product yields and are comparatively robust to metabolic stresses whereas lymphoid cells are more difficult to culture at an industrial scale. Given considerable cost of production, it is of utmost importance to maximize the yield of recombinant protein per bioreactor run. Choice of culture medium composition and bioreactor design and operation are parameters that impact yield and may be quite complex to optimize. More predictably, increases in the strength or transcriptional activity of the promoter controlling expression of product protein enhance yield. Incremental increases at the single cell level will translate into considerable improvements of product yield in high-density batch or fed-batch culture showing stationary phase gene expression at cell densities in the range of 10⁶ to 10⁷ cells/ml.

US5,866,359 describes a method of enhancing expression from an already strong hCMV promoter in CHO and NSO cells by co-expressiong adenoviral E1A protein from a weak promoter. E1A is a multifunctional transcription factor which may act on cell cycle regulation and has both independent transcriptional activating and repressing functional domains. The finetuning of E1A expression to appropriate low level expression is crucial for success of the co-expression approach in order to achieve the ideal balance in between gene transactivation whilst avoiding any negative impact on cell cycle progression.

As a disadvantage, apart from careful choice of the promoter driving E1A expression, this system blocks part of the protein synthesis capacity of the cell with E1A expression rather than expressing the recombinant protein of interest.

WO 95/17516 describes use of the murine immunoglobulin gamma 2A locus for targetting

10

15

25

30

an expression vector construct to a highly active gene locus in lymphoid cells of the B-cell lineage, e.g. widely used NS0 myeloma cells. NSO cells essentially are a tumor cell line of murine plasma or B-cells. Only in B-cells, the chromatin harboring the immunoglobulin loci is in its fully active, open state, allowing for high transcriptional activity of native immunoglobulin promoters or recombinant expression constructs integrated into those gene loci.

As a disadvantage, due to the principle of homologous recombination, the targetting sequence will target efficiently in murine cell lines only matching the sequence of the gamma 2 A targetting sequence harboring a recombinatorial hot spot; for high level expression, the gamma 2A locus region must be a transcriptionally active genomic region, limiting its effectiveness for homologous recombination to B-cell types.

It is an object of the present invention to devise another expression system for CHO protein expression in biotechnology which allows for enhanced expression from a standard promoter. According to the present invention, this aim is surprisingly achieved by equipping a gene expression vector for CHO cells with a gene targetting sequence having been originally devised for homologous recombination in murine B-cells.

- 20 Possible embodiments of the invention are shown in the figures. What is shown is:
 - Fig. 1 Relative expression levels of green fluorescent protein (GFP) from hCMV promoter and hCMV promoter in the presence of the IgG 2A hot spot sequence in transient transfection of CHO-K1 cells
 - Fig. 2 Relative GFP expression levels from hCMV promoter and hCMV promoter in the presence of the IgG 2A hot spot sequence in stably transfected CHO-K1 cells.
 - Fig. 3 Plasmid map of hCMV-MIE expression vector carrying IgG 2A targetting sequence

According to the present invention, a DNA sequence for expression of a recombinant gene in a mammalian cell comprises a recombinant product gene and a promoter for expressing the recombinant product gene, preferably a CMV promoter, and further comprises a

murine immunoglobulin gamma 2A locus DNA sequence or fragments or sequence variants thereof capable of enhancing expression from the promoter. According to the present invention, such a DNA sequence is useful expression vector construct for expression of recombinant product gene in CHO cells.

5

According to the present invention, the method of expressing a recombinant protein comprises the steps of

10

- a. culturing a CHO cell transfected with an expression vector comprising a promoter active in CHO cells driving expression of a recombinant product protein and further comprising the murine IgG 2 A gene locus DNA or a DNA sequence variant or DNA fragment thereof which is enhancing activity of said promoter, and
- b. harvesting the product protein

Immunoglobulin G (IgG) antibody.

A recombinant product gene according to the present invention is the product protein that 15

is sought to be expressed and harvested in high amount. It may be any protein of interest, e.g. therapeutic proteins such as interleukins or enzymes or subunits of multimeric proteins such as antibodies or fragments thereof. The recombinant product gene may include a signal sequence coding sequence portion allowing secretion of the once expressed polypeptide from the host producer cell. In a further prefered embodiment of the present invention, the product protein is a secreted protein. More preferably, the first or product protein is an antibody or engineered antibody or a fragment thereof, most preferably it is an

25

30

20

The DNA sequence of the murine immunoglobulin gamma 2A gene locus (IgG 2A) has originally been devised in WO 95/17516 for use as a genomic targetting sequence for generating stably recombinant lymphoid B-cell lines that show high expression of the recombinant gene product. B lymphocytes or plasma cells normally express extremely high levels of immunoglobulin RNA from the tle Ig heavy chain locus, probably due to celltype specific enhancer/transcription factor activity and open chromatin structure. The preferred murine immunoglobuline gamma 2A gene sequence of the present invention is the same as the targetting sequence used in WO 95/17516. It is a 5.1 kb BamHI genomic

20

25

fragment which includes all of the coding region of murine Ig gamma 2A except the most 5' part of the CH1 exon (Yamawaki-Kataoka, Y. et al., Proc. Natl. Acad. Sci. U.S.A. (1982) 79: 2623-2627; Hall, B. et al., Molecular Immunology (1989) 26:819-826; Yamawaki-Kataoka, Y. et al., Nucleic Acid Research (1981) 9: 1365-1381). According to the present invention, promotion of site-directed, homologous recombination is not the relevant property of the immunoglobulin gamma 2A gene sequence (IgG 2A). Accordingly, any sequence variant of said IgG 2A gene sequence or sequence fragment or variant sequence fragment that is functional in or capable of enhancing recombinant product gene expression from the promoter, preferably from a hCMV promoter as set forth below, both under condition of transient or stable expression in CHO cells is also encompassed by the present invention.

Such ,functional' variants encompass e.g. base insertions, deletions or point mutations and

be generated by methods well-known in the art, e.g. by primer-directed PCR, 'error-prone' PCR, 'gene-shuffling' termed PCR-reassembly of overlapping DNA fragments or by invivo random mutagenesis of bacterial clones followed by library transfection and functional selection in CHO cells. For instance, random mutagenesis can be achieved by alkylating chemicals or UV-irradiations as described in Miller, J., Experiments in Molecular Genetics, Cold Spring Harbor Laboratory 1972). Optionally, a natural mutator-strain of a host bacterium may be used.

Preferably, such variant sequence or sequence fragment is at least 65%, more preferably 75%, most preferably 90% homologous in DNA sequence to the corresponding part of the natural murine immunoglobuline gamma 2A gene locus. For instance, it is possible to insert a Sal I restriction site at the naturally occurring Stu I site present 39 bp upstream of membrane exon 2 (M2) to provide a unique site for linearization within the murine immunoglobulin gamma 2A sequence; such sequence variant was originally devised for site-specific recombination targetting, but can as well be employed in the context of the present invention.

A 'promoter' is defined as a DNA sequence that directs RNA polymerase to bind to DNA and intiates RNA synthesis. According to the present invention, it is a promoter that is active in CHO cells. Such a promoter preferably is a strong promoter. A strong promoter is one which causes mRNAs to be initated at high frequency equal to or higher than that of

present invention.

5

10

15

20

25

30

hCMV core promoter/enhancer fragment (described in US5168062) in CHO-KI cells. Such promoter may be a cell-type dependent strong promoter, as are cited in US5589392, or preferably is a ubiquitously active strong promoter, more preferably a constitutively active viral promoter such as e.g. early and late promoters of the SV40 virus, the immediate early promoter of the human cytomegalovirus (hCMV) or of murine cytomegalovirus (mCMV), the thymidine kinase promoter (TK) of Herpes Simplex virus or the Rous Sarcoma Virus long terminal repeat promoter (RS-LTR), more preferably it is the hCMV-MIE promoter as defined by the 2.1 kb Pst I fragment described in US 5,385,839 and/or EP-323 997-A1 or a functional part thereof having promoter activity. The hCMV promoter construct harboring the complete first functional intron of the major immediate early (MIE) gene of hCMV, as set forth in EP-323 997-A1, is a particularly preferred embodiment of the

Preferably a hCMV promoter employed in the present invention lacks the 'modulator' sequence part in the upstream/enhancer portion of the promoter. The 'modulator' sequence has been found to be detrimental to hCMV promoter activity in CHO cells and stretches from position -750 to position -1150 relative to the MIE transcription start site (Meier et al., 1996, Intervirology 39: 331-342, Regulation of hCMV immediate-early gene expression), in particular in transient transfection. Without the modulator sequence, the enhancing effect of the presence of the IgG 2A host spot sequence on (modulator negative or mod- for short) hCMV promoter is even more pronounced.

A transient transfection is characterised by non-appliance of any selection pressure for a vector borne selection marker. A pool or batch of cells originating from a transient transfection is a pooled cell population that comprises cells which have taken up and do express and cells that have not taken up the foreign DNA. Cells that express the foreign expression cassette do usually not have integrated the transfected DNA into their genome yet and tend to lose the foreign DNA and to overgrow transfected cells in the population upon culture of the transiently transfected cell pool. Therefore expression is strongest in the period immediately following transfection and decreases with time. Preferably, a transient transfectant according to the present invention is understood as a cell that is maintained in cell culture in the absence of selection pressure up to a time of 90 hours post transfection.

Preferably, a transfected CHO host cell according to the present invention is a stably transfected host cell, in particular in combination with a hCMV promoter as set forth above. Stable transfection means that newly introduced foreign DNA is becoming incorporated into genomic DNA, usually by random, non-homologous recombination 5 events; in case of a vector sequence, stable transfection according to the present invention may result in loss of vector sequence parts not directly related to expression of the recombinant product gene, such as e.g. bacterial copy number control regions rendered superfluous upon genomic integration. A transfected host cell has integrated at least part or different parts of the expression vector into the genome. Likewise, transfection of CHO 10 cells with two or several DNA fragments giving rise at least in vivo to functional equivalents of the essential elements of the expression vector of the invention, namely the product gene under control of a suitable promoter and the hot spot IgG 2A sequence, is contained in the definition of such transfected host cells. In vivo assembly of functional DNA sequences after transfection of fragmented DNA is described e.g. in WO 99/53046. It 15 is possible that such stable integration gives rise, upon exposure to further selection pressure for gene amplification, to double minute chromosomes in CHO cells. This is comprised in the present meaning of 'stable'. Upon random genomic integration of the expression vector of the present invention in CHO, the presence of the targetting sequence enhances promoter activity for expression of the recombinant product protein. Such effect 20 has not been observed nor could it have been anticipated upon homologous gene targetting in mature murine B-cell lines including plasmacytoma/myeloma cell lines; there, the IgG 2A targetting sequence served solely to increase the frequency of high-yielding homologous integrants since the IgG 2 A locus proved to be a recombinatorial 'hot spot'. As said before, the chromatin of the immunglobuline genomic region is in an open, highly 25 active state in suitably targetted B-cell lines.

'Expression vectors' are defined herein as DNA sequences that are required for transcription and the translation of their mRNAs in an appropriate mammalian host cell line after transfection with vector. An appropriately constructed expression vector should usually contain: at least one expressable marker selectable in animal cells, a limited number of useful restriction sites for insertion of the expression cassette for the recombinant product gene under control of an upstream promoter region. Where used in

വര് 🥞

5

10

15

20

25

30

particular for transient/episomal expression only, it may further comprise an origin of replication such as origin of Eppstein Barr Virus (EBV) or SV40 virus for autonomous replication/episomal maintenance in eukaryotic host cells but may be devoid of a selectable marker. Expression vectors are e.g., but are not limited to, linear DNA fragments, DNA fragments encompassing nuclear targetting sequences or are specially optimized for interaction with transfection reagents, animal viruses or suitable plasmids that can be shuttled and produced in bacteria. Any selection marker commonly employed such as thymidine kinase (tk), dihydrofolate reductase (DHFR) or glutamine synthetase (GS) may be used. In a preferred embodiment, an expressable GS selection marker is employed (Bebbington et al., 1992, High-level expression of a recombinant antibody from myeloma cells using a glutamine synthetase gene as an amplifiable selectable marker, Bio/Technology 10:169-175; Cockett et al., 1990, High level expression of tissue inhibitor of metalloproteinases in Chinese Hamster Ovary (CHO) cells using Glutamine synthetase gene amplification, Bio/Technology 8: 662-667). - The GS-system is one of only two systems that are of particular importance for the production of therapeutic proteins. In comparison to the dihydrofolate reductase (DHFR) system, the GS system offers a large time advantage during development because highly productive cell lines can often be created from the initial transectant thus avoiding the need for multiple rounds of selection in the presence of increasing concentrations of selective agent in order to achieve gene amplification (Brown et al., 1992, Process development for the production of recombinant antibodies using the glutamine synthetase (GS) system, Cytotechnology 9:231-236). It goes without saying that equivalent to a second transcription unit for expression of the marker gene, an expression unit could use a monocistronic expression cassette both for the product gene and the marker gene by employing e.g. internal ribosome entry sites as is routinely employed in the art. Vice versa, it goes without saying that the hot spot IgG 2 A sequence of the present invention and the expression cassette for the product protein comprising a promoter and/or marker cassette are not required to work in cis on a single expression vector; the elements can be well carried on separate co-transfected vectors or DNA fragments which may then be chromosomally integrated at a single, concatemeric integration site.

A further object of the present invention are CHO host cells transfected with the DNA sequences of the present invention. Further objects are a method for transfection of such m . 4)

5

host cells and a method for expression of the recombinant product gene in such host cells. The explanations and references made to preferred embodiments in the present specification of the invention relate likewise to all these further objects of the present invention. It is to be noted that a host cell transfected with the DNA sequence or vector of the present invention is to be construed as being a transiently or stably transfected cell line. Any transfection technique such as those well-known in the art, e.g. electoporation, Caphosphate precipitation, DEAE-dextrane transfection, lipofection, can be employed according to the present invention if appropriate for a given host cell type.

A suitable host cell line can be any chinese hamster ovary (CHO) cell line (Puck et al., 10 1958, J. Exp. Med. 108: 945-955). The term 'host cell' refers to cells capable of growth in culture and expressing a desired protein recombinant product protein. Suitable cell lines can be e.g. CHO K1 (ATCC CCL-61), CHO pro3-, CHO DG44, CHO P12 or the dhfr-CHO cell line DUK-BII (Chassin et al., PNAS 77, 1980, 4216-4220) or DUXB11 (Simonsen et al., PNAS 80, 1983, 2495-2499). In CHO cells, the immunoglobuline gene 15 loci are inactive and the chromatin is therefore in a densely packaged or closed state. Thus, any gene construct integrated in the immunoglobuline loci could not give rise to high-level expression of recombinant protein due to the specific state of chromatin, unless it would itself comprise flanking locus control regions promoting opening of the chromatin on both sides of the expression cassette. Further, immunoglobuline gene sequence, and in 20 particular the intron portions of it, show considerably divergence amongst species, e.g. from mouse to hamster. The promoter or enhancer elements of immunoglobline loci are both species and tissue specific and should be active in B-cells only. The murine IgG 2A sequence of the present invention enhances gene expression in CHO cells also in the absence of any natural immunoglobuline promoter that is giving rise to full-length 25 transcripts coding for complete IgG heavy chain. Preferably, the IgG 2A sequence of the present invention is devoid of such promoter. Surprisingly, the murine IgG 2 A targetting sequence even improved gene expression in CHO cells upon transient transfection of CHO cells with expression vectors according to the present invention (Fig. 1); such transient expression is a further preferred embodiment of a method according to the present 30 invention. In transient expression assays which are commonly taking place about 20-50 hours post transfection, the transfected vectors are maintained as episomal elements and are not yet integrated into the genome.

10

15

9

Suitable media and culture methods for mammalian cell lines are well-known in the art, as described in US 5633162 for instance. Examples of standard cell culture media for laboratory flask or low density cell culture and being adapted to the needs of particular cell types are for instance: Roswell Park Memorial Institute (RPMI) 1640 medium (Morre, G., The Journal of the American Medical Association, 199, p.519 f. 1967), L-15 medium (Leibovitz, A. et al., Amer. J. of Hygiene, 78, 1p.173 ff, 1963), Dulbecco's modified Eagle's medium (DMEM), Eagle's minimal essential medium (MEM), Ham's F12 medium (Ham, R. et al., Proc. Natl. Acad. Sc.53, p288 ff. 1965) or Iscoves' modified DMEM lacking albumin, transferrin and lecithin (Iscoves et al., J. Exp. med. 1, p. 923 ff., 1978). For instance, Ham's F10 or F12 media were specially designed for CHO cell culture. Other media specially adapted to CHO cell culture are described in EP-481 791. It is known that such culture media can be supplemented with fetal bovine serum (FBS, also called fetal calf serum FCS), the latter providing a natural source of a plethora of hormones and growth factors. The cell culture of mammalian cells is nowadays a routine operation well-described in scientific textbooks and manuals, it is covered in detail e.g. in R. Ian Fresney, Culture of Animal cells, a manual, 4th edition, Wiley-Liss/N.Y., 2000.

Preferably, the cell culture medium according to the present invention is devoid of fetal calf serum (FCS or FBS), which then is being termed 'serum-free'. Cells in serum-free 20 medium generally require insulin and transferrin in a serum-free medium for optimal growth. Transferrin may at least partially be substituted by non-peptide chelating agents or siderophores such as tropolone as described in WO 94/02592 or increased levels of a source of anorganic iron favorably in conjunction with antioxidants such as vitamin C. Most cell lines require one or more of synthetic growth factors (comprising recombinant 25 polypeptides), including e.g. epidermal growth factor (EGF), fibroblast growth factor (FGF), insulin like growth factors I and II (IGFI, IGFII), etc.. Other classes of factors which may be necessary include: prostaglandins, transport and binding proteins (e.g. ceruloplasmin, high and low density lipoproteins, bovine serum albumin (BSA)), hormones, including steroid-hormones, and fatty acids. Polypeptide factor testing is best 30 done in a stepwise fashion testing new polypeptide factors in the presence of those found to be growth stimulatory. Those growth factors are synthetic or recombinant. There a several methodological approaches well-known in animal cell culture, an exemplary being

10

15

20

25

30

described in the following. The initial step is to obtain conditions where the cells will survive and/or grow slowly for 3-6 days after transfer from serum-supplemented culture medium. In most cell types, this is at least in part a function of inoculum density. Once the optimal hormone/growth factor/polypeptide supplement is found, the inoculum density required for survival will decrease. In a more preferred embodiment, the cell culture medium is protein-free, that is free both of fetal serum and individual protein growth factor supplements or other protein such as recombinant transferrin.

A possible embodiment of one method of the present invention, namely expression and harvest of the recombinant product protein, is high-density growth of the animal host cells e.g. in an industrial fed-batch bioreactor. Conventional downstream processing may then be applied. Consequently, a high-density growth culture medium has to be employed. Such high-density growth media can usually be supplemented with nutrients such as all amino acids, energy sources such as glucose in the range given above, inorganic salts, vitamins, trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), buffers, the four nucleosides or their corresponding nucleotides, antioxidants such as Glutathione (reduced), Vitamine C and other components such as important membrane lipids, e.g. cholesterol or phosphatidylcholine or lipid precursors, e.g. choline or inositol. A high-density medium will be enriched in most or all of these compounds, and will, except for the inorganic salts based on which the osmolarity of the essentially isotonic medium is regulated, comprise them in higher amounts (fortified) than the afore mentioned standard media as can be incurred from GB2251 249 in comparison with RPMI 1640. Preferably, a high-density culture medium according to the present invention is balancedly fortified in that all amino acids except for Tryptophane are in excess of 75 mg/l culture medium. Preferably, in conjunction with the general amino acid requirement, Glutamine and/or Asparagine are in excess of 1 g/l, more preferably of 2 g/l of high-density culture medium. In the context of the present invention, high-density cell culture is defined as a population of animal cells having temporarily a density of viable cells of at least or in excess of 10⁵ cells/ml, preferably of at least or in excess of 10⁶ cells/ml, and which population has been continously grown from a single cell or inoculum of lower viable cell density in a cell culture medium in a constant or increasing culture volume.

10

15

20

25

30

In a further prefered embodiment, the fed-batch culture is a culture system wherein at least Glutamine, optionally with one or several other amino acids, preferably glycine, is fed to the cell culture as described in GB2251249 for maintaing their concentration in the medium, apart from controlling glucose concentration by spearate feed. More preferably, the feed of glutamine and optionally one or several other amino acids is combined with feeding one or more energy sources such as glucose to the cell culture as described in EP-229 809-A. Feed is usually initiated at 25-60 hours after start of the culture; for instance, it is useful to start feed when cells have reached a density of about 10⁶ cells/ml. It is well known in the art that in cultured animal cells, 'glutaminolysis' (McKeehan et al., 1984, Glutaminolysis in animal cells,in: Carbohydrate Metabolism in Cultured Cells, ed. M.J. Morgan, Plenum Press, New York, pp. 11-150) may become an important source of energy during growth phase. The total glutamine and/or asparagine feed (for substitution of glutamine by asparagine, see Kurano, N. et al., 1990, J. Biotechnology 15, 113-128) is usually in the range from 0.5 to 10 g per l, preferably from 1 to 2 g per l culture volume; other amino acids that can be present in the feed are from 10 to 300 mg total feed per litre of culture, in particular glycine, lysine, arginine, valine, isoleucine and leucine are usually fed at higher amounts of at least 150 to 200 mg as compared to the other amino acids. The feed can be added as shot-addition or as contionusly pumped feed, preferably the feed is almost continously pumped into the bioreactor. It goes without saying that the pH is carefully controlled during fed-batch cultivation in a bioreactor at an approximately physiological pH optimal for a given cell line by addition of base or buffer. When glucose is used as an energy source the total glucose feed is usually from 1 to 10, preferably from 3 to 6 grams per litre of the culture. Apart from inclusion of amino acids, the feed preferably comprises a low amount of choline in the range of 5 to 20 mg per litre of culture. More preferably, such feed of choline is combined with supplementation of ethanolamine essentially as described in US 6048728, in particular in combination with feeding glutamine. It goes without saying that upon use of the GS-marker system, lower amounts of glutamine will be required as compared to a non-GS expression system since accumulation of excessive glutamine in addition to the endogenously produced would give rise to ammonia production and concomittant toxicity. For GS, glutamine in the medium or feed is mostly substituted by its equivalents and/or precursors, that is asparagine and/or glutamate.

10

15

It is a further, independent object of the present invention to devise an expression vector comprising at least a (first) transcription unit for a product gene, giving rise to product protein upon expression in a host cell, and which transcription unit is under the control of the mouse Cytomegalovirus promoter (mCMV promoter), and further comprising a second transcription unit comprising a glutamine synthetase (GS) marker gene. Such a product gene, or gene of interest (GOI) as it may be termed, can be e.g. an immunoglobulin coding sequence. A glutamine synthetase marker gene is any enzymatically active GS coding sequence, be it a natural gene sequence or a variant thereof. The above definitions of 'functional variant' as set forth above apply here as well including the preferred ranges of sequence homology. Preferably, the GS marker gene is a mammalian GS marker gene or derived thereof. Surprisingly, such expression vector allows for much higher transfection rates upon transfection in CHO cells than does e.g. an expression vector in which the first transcription unit harboring the gene of interest is under control of the hCMV promoter. This despite the fact that in CHO cells, transcriptional activity of the mCMV promoter is much higher than that of hCMV promoter; usually it is believed that upon transfection, higher metabolic load reduces clonal survival upon transfection, resulting in lower numbers of transfectants. Thus the effect can not be correlated in an obvious manner with the amount or unexpected toxicity of product protein expressed, the latter possibly adversely affecting growth of transfectants. Indeed, the finding is the very opposite of any 20 expectation of a skilled person. Further objects according to the present invention are animal host cells, in particular CHO cells, transfected with such an expression vector which vector can be maintained episomally or can be stably integrated in the genome and a respective transfection method. Likewise, transfection of animal cells, in particular CHO cells, with two or more gene 25 fragments giving rise in-vivo to functional equivalents of the transcription units of the present object of the invention, is within the definition of such transfected host cells. Preferably, said host cells are stably transfected cells, meaning that the first and second

30

A further object is the use of mCMV promoter to enhance transfection rate in CHO cells, preferably when using an expression vector comprising at least a first transcription unit for a product gene which first unit is giving rise to product protein upon expression in a host

transcription unit are chromosomally integrated.

10

15

20

25

30

cell and which first transcription unit is further under the control of the mouse Cytomegalovirus promoter (mCMV promoter), and further comprising a second transcription unit comprising a glutamine synthetase (GS) marker gene. It may also be possible to transfect the first and second expression borne on different vectors, or as isolated gene fragments harboring individual expression units. Further, it may be possible to transfect a CHO cell that is already recombinant for and expresses GS with a first transcription unit harboring mCMV. According to the present invention, 'enhancing transfection rate' is defining by comparing transfection rate in the presence of the mCMV promoter and expression vector according to the present invention with the transfection rate of the same expression vector and host cell under identical transfection and cell culture conditions except that in the expression vector, the mCMV promoter is substituted to the hCMV-first intron enhancer/promoter construct as defined in US 5658 759 and as set forth e.g. in sequence ID. No. 3 of the present invention. This hCMV-intron MIE-promoter construct, for a given identical product gene, serves as a standard for determining the claimed effect of enhanced transfection rates. Preferably, use of mCMV promoter results in

All relevant definitions given further above apply likewise to the present, independent objects of the invention. It must be stressed that the present object of the invention does not require the presence of the murine IgG 2 A targetting sequence as a prerequisite.

at least 10-times enhanced transfection rate.

Murine cytomegalovirus (mCMV) is a member of the highly diverse group of herpesviridae. Even amongst cytomegaloviruses of different host species there can be wide variation. For example, mCMV differs considerably from the human cytomegalovirus (hCMV) with respect to biological properties, immediate early (IE) gene organization, and overall nucleotide sequence. The 235-kbp genome of mCMV also lacks large internal and terminal repeat characteristics of hCMV. Accordingly, no isomeric forms of the MCMV genome exist (Ebeling, A. et al., (1983), J. Virol. 47, 421-433; Mercer, J. A. et al., (1983), Virology 129, 94-106). According to the present invention, it is possible to employ the promoter region essentially corresponding to a large approx. 2.1 kb PstI fragment described in US 4968615 or any functional fragment thereof. In a more preferred embodiment, the mCMV promoter fragment employed comprises the transcription start site (+0) and extends upstream to about position –500. Surprisingly, such fragment has

5

10

15

20

25

30

been found to promote stronger expression than a promoter cassette extending 800 bp further upstream beyond position -500. In a most preferred embodiment, a core promoter region is employed that extends from the transcription start site upstream but to the Xho I restriction site at about position -150 from the natural transcription start site or even extending but to position -100 upstream from the natural transcriptions start site. It goes without saying that the transcription start site might be engineered in order to comprise a suitable restriction site for insertion of the recombinant product gene.

According to the present invention, it is also possible that the first transcription unit that is under control of the mCMV promoter harbors at least one intron sequence. Such measure is well-known in the art for stabilising RNA transcripts and for promoting efficient protein synthesis from the corresponding mRNA. For efficient protein synthesis without having regard to the claimed effect on transfection rate, it is however not advisable to include the first, natural intron of mCMV in the mCMV promoter construct. In contrast to the situation with hCMV promoter (cf. US 5591639), such natural first intron of mCMV was found to decrease expression of a recombinant gene from the mCMV promoter and is therefore excluded in a further preferred embodiment.

Examples of preferred, possible embodiments of GS marker gene cassettes are given in the sequence listings. Seq IDs No. 1 (pEE 15.1 hCMV/GFP + hot spot) +2 (pEE 14.4 hCMV/GFP) give examples of suitable GS-gene cassettes that are expressed from the SV40 (early and late, respectively) promoter, a weak to medium level promoter, further comprising an expression cassette for GFP (Green fluorescent protein) that is under control of the hCMV promoter. Seq. ID No. 1 describes a GS cDNA sequence described in more detail in the figure legend of Fig. 3, under control of the SV40 early promoter. Seq. ID No. 2 specifies an artifical GS-minigene cassette comprising an intron that is under control of the SV40 late promoter. CHO cells are not naturally glutamine auxothropic, therefore selection schemes as e.g. described in Cockett et al., 1990, High level expression of tissue inhibitor of metalloproteinases in Chinese Hamster Ovary (CHO) cells using Glutamine synthetase gene amplification, Bio/Technology 8: 662-667, can be applied. Examples of suitable transfection methods for CHO cells are equally given therein; it is possible to employ e.g. classic calcium phosphate precipitation or more modern lipofection

techniques. Transfection rate is routinely defined as the number of positively transfected cells (transient transfection) or clones (stable transfection after selection period) obtained from a pool of cells subjected to transfection. The purported effect of the present object of invention can be seen e.g. by transfecting CHO-K1 cells by lipofection (any commercial reagent and manufacturers protocol) with the plasmids of either Seq. ID No. 3 (pEE 12.4 hCMV-GFP + SV40 early promoter/GS cDNA) or Seq. ID No. 4 (pEE 12.4 mCMV-GFP + SV40 early promoter/GS cDNA). Transfected cells may be grown in any conventional culture medium. The culture medium may be a fetal serum-supplemented or serum-free medium as has been defined above. Preferably, the cell culture medium is a serum-supplemented medium, more preferably a cell culture medium that has been supplemented with at least 1% (v/v) fetal serum, most preferably with at least 5% (v/v) fetal serum such as fetal calf serum or fetal bovine serum. In another preferred embodiment, the transfection method carried out is electroporation.

10

5

Experiments

Experiment 1

5

10

15

30

<u>Transient and stable expression of GFP vector comprising hot spot sequence in CHO-K1</u> cells

16

CHO-K1 cells (ATCC CCL-61) were adapted and cultured in normal cell culture medium GMEM –S (Gibco, UK) with 10% FCS. - For GS selection, the medium must be completely free of glutamine as set forth in table 1 below; this necessitates use of dialysed FCS. - All culturing was carried out in shake flask at 36.5 °C with orbital shaking at 125 rpm. Lipofectin (Superfectin™, Gibco, UK) was used for transfection and green fluorescence of transfectant pool was measured in a FACS with excitation at 488 nm. For every GS/GFP vector construct, transfection was carried out independently five times, all data being the average from five independently analyzed pools. Starting with transient transfectants 48 h post-transfection, the top scoring 10% highly expressing cells of the viable cell pool in the cell count vs. fluorescence diagram were selected to determine mean fluorescence (Fig.1). Viable cell population has been preselected by gating in the Forward vs. sideward scatter diagram.

For generating stable transfectants, GS marker was selected 24 hours post-transfection by supplementing the glutamine-free medium with 25 μM MSX (methionine sulphoximine, Crockett et al., ibd.) and continuing cell culture with regular splitting of cultures for 26 days. Note the impact of medium levels of other amino acids on the potency of MSX for selection, see Bebbington et al., US 5827 739. Flurorescence analysis was then performed again as outlined above (Fig. 2).

Untransfected cells served as negative control. The hot spot vector (pEE 15.1 'hCMV + hot spot') driving expression of GFP under control of the hCMV promoter comprising the first complete intron of CMV is given in Seq. ID No. 1 and essentially is the pEE 15.1 vector shown in Fig. 3 into which the GFP sequence was inserted into the Eco RI restriction site in the polylinker. pEE 12.4 'hCMV' corresponding to Seq. ID No. 3 is identical to pEE 15.1 'hCMV + hot spot' except that it does not comprise the 5.1 kb Bam H1 fragment harboring the IgG 2A sequence. pEE 12.4 served as a vector control. A further vector

5

10

15

20

25

30

control pEE 12.4 'hCMV(Kozak-)' was generated by mutating the Kozak sequence of the cloning site coninciding with the translation start site (GCCGCCACCATGG) to a frameshifted functional Kozak sequence that (ACCATGGTCCATGG) by primer directed mutagensis (Sambrook et al., Molecular cloning, Cold Spring Harbor 1983), attentuating the original Kozak and translation start site. The vector of Seq. ID No. 1 was further engineered to delete the 400 bp modulator region of hCMV enhancer portion, deleting the enhancer elements upstream of -750 from the transcription start site, giving rise to pEE 15.1 'hCMV(mod-)/GS cDNA'. By exchange of the GScDNA cassette of pEE 15.1 (s. Fig. 3) with the GS minigene of pEE 14.4 'hCMV(mod-)' /GFP, corresponding to Seq. ID No. 2, the vector pEE 15.1 'hCMV(mod-)/GS minigene' was created. Thus all transfected cells harbored a plasmid vector comprising the GFP coding sequence. The GS minigene contains a single, first intron of the GS gene and about 1 kb of 3' flanking DNA under the control of the SV40 late promoter; the 3' part of the genomic GS DNA is believed to cause a higher copy-number of vector DNA and thus of GS in transfected cells (see US4770359, Bebbington et al.). Whereas all hCMV vectors employed in the present study express the GS maker gene from its cDNA sequence, use of the GS minigene was included as a further control in order to exclude potential effects of GS copy number and expression level.

For generation and expression analysis of stably transfected CHO cells, transfections were performed with linearized hot spot vector pEE 15.1 'hCMV+ hot spot' vector. Sal I linearized plasmid was cut in the IgG 2A comprising sequence portion, free DNA ends potentially stimulating recombination with genomic regions sharing a certain degree of homology with the flanking DNA portions, testing for potential targetting effects of murine IgG 2A in hamster CHO cells. Pvu I cut in the bacterial lactamase marker gene and therefore could promote but heterologous random recombination. Indeed, the mean fluorescence was higher in the Pvu I linearized transfectants showing both some influence of vector linearization as well as that targetting to immunoglobuline loci in CHO cells may not account for the effect of the present invention. In addition, the effect of enhanced promoter activity was consistingly observed in transiently transfected cell populations, nicely correlating with relative strength of individual vector constructs. Clearly, genomic integration is not involved at this early stage of transfection.

Fig. 3 shows vector pEE 15.1 of approximately 12 830 bp. A detailed description of the GS

marker and the hCMV-p/intron expression cassette can be found in US5827739 and US5591639, pEE 15.1 is a possible embodiment of an expression vector according to the present invention, except that the DNA sequence coding for the recombinant product protein has not yet been inserted in the polylinker site. The complete 13535 bp sequence of the pEE15.1 construct harboring GFP is given in Seq. ID No. 1: Therein, the GFP coding sequence was inserted in-frame in the Eco R I restriction site centered at base position 12 814; the introduction of the unique restriction site harboring the ATG start codon and optimizing the Kozak sequence environment of the start codon is described in detail in US 5591639. Thus, the expression of GFP protein is under control of the hCMV-major immediate early gene promoter (hCMV-MIE or hCMV for short) immediately followed by the first intron of hCMV-MIE gene followed by the Nco I site (s. US 5591 639). Polyadenlyation is ensured by the SV40 poly A site further downstream of the polylinker insertion site. pEE 15.1 further harbors a cDNA sequence coding for glutamine synthetase (GS) from hamster that is under control of the SV 40 early promoter and is followed by an SV40 intron +poly A sequence. The IgG 2A gene locus or 'hot spot' sequence (hatched boxes CH1, Hi, CH2, CH3, M1, M2 standing for Heavy chain constant region, hinge, membrane anchor) is the 5.1 kb BamH I fragment of the murine IgG 2 A locus already described in WO 9517516 and the references cited therein. Unique restriction sites Pvu I and Sal I are shown.

20

25

30

5

10

15

Experiment 2

Electroporation of CHO cells with mCMV p12.4 -GFP construct (Seq. ID No.4)

Attached CHO-K1 cells (ATCC CCL-61) were cultured in Iscoves' DMEM medium essentially as described in EP-481 791 comprising 2 mM Glutamine which was further supplemented with 10% FCS. Optionally, the G-MEM medium stated in table 1 and further comprising 2 mM Glutamine could be used prior to GS marker selection as in experiment 1. The cells were detached, pelleted and resuspended twice in serum-free medium, finally at a density of 5.3 x10⁶ cells/ml. Per 750 μl electroporation batch, a total of 4x10⁶ cells was electroporated. Electroporation was carried out as described in Methods in Molecular Biology, ed. JA Nickoloff ed, Humana Press 1995, Vol. 48/Chap. 8: Animal cell electroporation and electrofusion protocols. p12.4 mCMV-GFP vector DNA (sequence ID No. 4) was linearized. 50μl (20μg) DNA were added to 750μl cells in electroporation

20

25

30

35

cuvette and electroporate - 300 Volts / 750 μ Fd - expecting an electorporation time of around 12 - 14 msec. Following electroporation 800 μ l volume of cells was transferred into 25 ml of modified Glasgow-MEM (GMEM, Gibco) culture medium for GS selection (comprising 10% fetal serum but no glutamine, for details see table 1) in a T75 flask.

Divide into 2 x T75 flasks by moving 12.9 mls into a second flask and incubate overnight at 37°C in 10% CO₂

On the next day 37.5ml of GS-selection GMEM culture medium supplemented with 10% FBS + 33.3µM MSX (methionine sulphoximine) were added. Thus MSX was finally ~25µM. Transfectants were counted after further incubation for 26 days by colony count per flask. Upon microscopic inspection in a standard inverted microscope for inspection of culture flasks, positive colonies brightly lit up in light green and could be easily counted.

The mCMV construct of Seq. ID No. 4 yielded up to 20 times more foci than did cells that were transfected in parallel with the hCMV construct of Seq. ID No. 3. The vector constructs only differed in the CMV promoter elements driving GFP expression, the remaining vector parts of the vectors were identical (including GS-marker; cDNA GS-marker cassette of p12.4). If cells were diluted out into 96 well plates immediately following transfection, many more colonies come up from mCMV transfected cells (>400 colonies) than from hCMV transfected cells (about 45 colonies).

TABLE 1: Medium for GS selection

A. Stock Solutions

- 1. Double distilled water autoclaved in 400 ml aliquots
- 2. 10 x Glasgow MEM (GMEM) without glutamine (GIBCO: 042-2541 in UK). Store at 4°C.
- 3. 7.5% sodium bicarbonate (GIBCO: 043-05080 in UK; 670-5080 in US). Store at 4°C.
- 4. 100 x non-essential amino acids (NEAA) (GIBCO: 043-01140 in UK; 320-1140 in US). Store at 4°C.
- 100 x Glutamate + Asparagine (G + A): add 600 mg glutamic acid and 600 mg asparagines (Sigma).
 Make up to 100 ml in distilled water and sterilize by passing through a sterile 2μm filter (Nalgene).
 Store at 4°C.
- 100mM sodium pyruvate (GIBCO: 043-01360 in UK; 320-1360 in US)

10

15

7.	50 x nucleosides:	35 mg adenosine
		35 mg guanosine
		35 mg cytidine
•		35 mg uridine
		12 mg thymidine

(each from Sigma). Make up to 100ml with water, filter sterilise and store at -20°C in 10ml aliquots.

- 8. Dialysed FCS (GIBCO: 014-06300). Heat inactivate at 56°C for 30 min and store at -20°C. It is essential to use dialysed FCS when using GS selection.
 - 9. Penicilin-streptomycin at 5000 units/ml (P/S: GIBCO: 043-05070 in UK; 600-5070 in US).
 - 10. 100 mM L.MSX (Sigma): prepare 18 mg/ml solution in PBS. Filter sterilise and store at -20°C.

B.Medium Preparation

Add the following in the order given using aseptic technique to make GMEM-S medium

20	1. Water	400 ml
	2. 10 x GMEM	50 ml
	3. Sodium bicarbonate	18.1 ml
	4. NEAA	5 ml
	5. G + A	5 ml
25	6. Sodium pyruvate	5 ml
	7. Nucleosides	10 ml
	8. Dialysed FCS	50 ml
	9. Penicillin-streptomycin	5 ml

GMEM-S contains the non-essential amino acids, alanine, aspartate, glycine, proline and serine (100 μM), glutamate and asparagines (500 μM), and adenosine, guanosine, cytidine and uridine (30 μM), and thymidine (10 μM).

Claims

5


20


25

- 1. CHO cell transfected with an expression vector comprising a promoter that is active in CHO cells and that is driving expression of a recombinant product protein and further comprising a portion from the murine IgG 2 A gene locus DNA which portion is enhancing activity of said promoter.
- 2. CHO cell according to claim 1, characterized in that the vector further comprises a transcription unit encoding a selectable marker, preferably a glutamin synthetase (GS) marker.
 - 3. CHO cell according to claim 1 or 2, characterized in the CHO cell is stably transfected.
- 4. Method of expressing a recombinant protein, comprising the steps of
 - c. culturing a CHO cell transfected with an expression vector comprising a promoter active in CHO cells driving expression of a recombinant product protein and further comprising the murine IgG 2 A gene locus DNA or a DNA sequence variant or DNA fragment thereof which is enhancing activity of said promoter, and
 - d. harvesting the product protein
 - 5. Method according to claim 4, characterised in that the promoter is a strong viral promoter, preferably the hCMV promoter.
 - 6. Method according to one of claims 4 or 5, characterised in that the IgG 2A gene locus portion does lack the natural immunoglobulin promoter.
- 7. Mammalian expression vector comprising at least a first transcription unit for a product gene which transcription unit is under the control of the mCMV promoter, and further comprising a second transcription unit comprising a glutamine synthetase (GS) marker gene.
 - 8. CHO cell transfected with the vector of claim 7.

PCT/EP2003/007946


1/3





3/3

Fig. 3



540



LBP30AND31.ST25.txt SEQUENCE LISTING

Lonza Biologics plc. <110> Method of expressing recombinant protein in CHO cells <120> <130> LBP30+31 <140> NA 25/01 <141> 2002-07-. <160> PatentIn version 3.1 <170> <210> 1 <211> 6679 <212> DNA <213> Hamster sp. <220> <221> misc_feature Seq. ID. No. 4: circular plasmid GS vector p12.4 short mCMV-GFP / <223> clone 3 <400> 1 gaattcattg atcataatca gccataccac atttgtagag gttttacttg ctttaaaaaa 60 cctcccacac ctcccctga acctgaaaca taaaatgaat gcaattgttg ttgttaactt 120 gtttattgca gcttataatg gttacaaata aagcaatagc atcacaaatt tcacaaataa 180 agcatttttt tcactgcatt ctagttgtgg tttgtccaaa ctcatcaatg tatcttatca 240 tgtctggcgg ccgcgacctg caggcgcaga actggtaggt atggaagatc cctcgagatc 300 cattgtgctg gcggtaggcg agcagcgcct gcctgaagct gcgggcattc ccagtcagaa 360 atgagcgcca gtcgtcgtcg gctctcggca ccgaagtgct atgattctcc gccagcatgg 420

cttcggccag tgcgtcgagc agcgcccgct tgttcctgaa gtgccagtaa agcgccggct

gctgaacccc caaccgttcc gccagtttgc gtgtcgtcag accgtctacg ccgacctcgt



tcaacaggtc	cagggcggca	cggatcactg	tattcggctg	caactttgtc	atgcttgaca	600
				gtgataaaga		660
				atattcgcgg		720
				accgatgggg		780
				atggtggcag		840
cgggggactg	ttgggcgcca	tctccttgca	tgcaccattc	cttgcggcgg	cggtgctcaa	900
cggcctcaac	ctactactgg	gctgcttcct	aatgcaggag	tcgcataagg	gagagcgtcg	960
acctcgggcc	gcgttgctgg	cgtttttcca	taggctccgc	cccctgacg	agcatcacaa	1020
aaatcgacgc	tcaagtcaga	ggtggcgaaa	cccgacagga	ctataaagat	accaggcgtt	1080
tcccctgga	agctccctcg	tgcgctctcc	tgttccgacc	ctgccgctta	ccggatacct	1140
gtccgccttt	ctcccttcgg	gaagcgtggc	gctttctcat	agctcacgct	gtaggtatct	1200
cagttcggtg	taggtcgttc	gctccaagct	gggctgtgtg	cacgaacccc	ccgttcagcc	1260
cgaccgctgc	gccttatccg	gtaactatcg	tcttgagtcc	aacccggtaa	gacacgactt	1320
atcgccactg	gcagcagcca	ctggtaacag	gattagcaga	gcgaggtatg	taggcggtgc	1380
tacagagttc	ttgaagtggt	ggcctaacta	cggctacact	agaagaacag	tatttggtat	1440
ctgcgctctg	ctgaagccag	ttaccttcgg	aaaaagagtt	ggtagctctt	gatccggcaa	1500
acaaaccacc	gctggtagcg	gtggttttt	tgtttgcaag	cagcagatta	cgcgcagaaa	1560
aaaaggatct	caagaagatc	ctttgatctt	ttctacgggg	tctgacgctc	agtggaacga	1620
aaactcacgt	taagggattt	tggtcatgag	attatcaaaa	aggatcttca	cctagatcct	1680
tttaaattaa	aaatgaagtt	ttaaatcaat	ctaaagtata	tatgagtaaa	cttggtctga	1740
cagttaccaa	tgcttaatca	gtgaggcacc	tatctcagcg	atctgtctat	ttcgttcatc	1800
catagttgcc	tgactccccg	tcgtgtagat	aactacgata	cgggagggct	taccatctgg	1860
ccccagtgct	gcaatgatac	cgcgagaccc	acgctcaccg	gctccagatt	tatcagcaat	1920
aaaccagcca	gccggaaggg	ccgagcgcag	aagtggtcct	gcaactttat	ccgcctccat	1980
ccagtctatt	aattgttgcc	gggaagctag	agtaagtagt	tcgccagtta	atagtttgcg	2040
caacgttgtt	gccattgcta	caggcatcgt	ggtgtcacgc	tcgtcgtttg	gtatggcttc	2100
attcagctcc	ggttcccaac	gatcaaggcg	agttacatga	tcccccatgt	tgtgcaaaaa	2160
agcggttagc	tccttcggtc	ctccgatcgt	tgtcagaagt	aagttggccg	cagtgttatc	2220
actcatggtt	atggcagcac	tgcataattc	tcttactgtc	atgccatccg	taagatgctt	2280
ttctgtgact	ggtgagtact	caaccaagtc	attctgagaa	tagtgtatgc	ggcgaccgag	2340
ttgctcttgc	ccggcgtcaa	tacgggataa	taccgcgcca	catagcagaa	ctttaaaagt	2400
gctcatcatt	ggaaaacgtt	cttcggggcg	aaaactctca	aggatcttac	cgctgttgag	2460
atccagttcg	atgtaaccca	ctcgtgcacc	caactgatct	tcagcatctt	ttactttcac	2520
cagcgtttct	gggtgagcaa	aaacaggaag	gcaaaatgcc	gcaaaaaagg	gaataagggc	2580

LBP30AND31.ST25.txt gacacggaaa tgttgaatac tcatactctt cctttttcaa tattattgaa gcatttatca	2640
gggttattgt ctcatgagcg gatacatatt tgaatgtatt tagaaaaata aacaaatagg	2700
ggttccgcgc acatttcccc gaaaagtgcc acctgacgtc taagaaacca ttattatcat	2760
gacattaacc tataaaaata ggcgtatcac gaggccctga tggctctttg cggcacccat	2820
cgttcgtaat gttccgtggc accgaggaca accctcaaga gaaaatgtaa tcacactggc	2880
tcaccttcgg gtgggccttt ctgcgtttat aaggagacac tttatgttta agaaggttgg	2940
taaattcctt gcggctttgg cagccaagct agatccggct gtggaatgtg tgtcagttag	3000
ggtgtggaaa gtccccaggc tccccagcag gcagaagtat gcaaagcatg catctcaatt	3060
agtcagcaac caggtgtgga aagtccccag gctccccagc aggcagaagt atgcaaagca	3120
tgcatctcaa ttagtcagca accatagtcc cgcccctaac tccgcccatc ccgcccctaa	3180
ctccgcccag ttccgcccat tctccgcccc atggctgact aattttttt atttatgcag	3240
aggccgaggc cgcctcggcc tctgagctat tccagaagta gtgaggaggc ttttttggag	3300
gcctaggctt ttgcaaaaag ctagcttggg gccaccgctc agagcacctt ccaccatggc	3360
cacctcagca agttcccact tgaacaaaaa catcaagcaa atgtacttgt gcctgcccca	3420
gggtgagaaa gtccaagcca tgtatatctg ggttgatggt actggagaag gactgcgctg	3480
caaaacccgc accctggact gtgagcccaa gtgtgtagaa gagttacctg agtggaattt	3540
tgatggctct agtacctttc agtctgaggg ctccaacagt gacatgtatc tcagccctgt	3600
tgccatgttt cgggacccct tccgcagaga tcccaacaag ctggtgttct gtgaagtttt	3660
caagtacaac cggaagcctg cagagaccaa tttaaggcac tcgtgtaaac ggataatgga	3720
catggtgagc aaccagcacc cctggtttgg aatggaacag gagtatactc tgatgggaac	3780
agatgggcac ccttttggtt ggccttccaa tggctttcct gggccccaag gtccgtatta	3840
ctgtggtgtg ggcgcagaca aagcctatgg cagggatatc gtggaggctc actaccgcgc	3900
ctgcttgtat gctggggtca agattacagg aacaaatgct gaggtcatgc ctgcccagtg	3960
ggaactccaa ataggaccct gtgaaggaat ccgcatggga gatcatctct gggtggcccg	4020
tttcatcttg catcgagtat gtgaagactt tggggtaata gcaacctttg accccaagcc	4080
cattcctggg aactggaatg gtgcaggctg ccataccaac tttagcacca aggccatgcg	4140
ggaggagaat ggtctgaagc acatcgagga ggccatcgag aaactaagca agcggcaccg	4200
gtaccacatt cgagcctacg atcccaaggg gggcctggac aatgcccgtg gtctgactgg	4260
gttccacgaa acgtccaaca tcaacgactt ttctgctggt gtcgccaatc gcagtgccag	4320
catccgcatt ccccggactg tcggccagga gaagaaaggt tactttgaag accgcggccc	4380
ctctgccaat tgtgacccct ttgcagtgac agaagccatc gtccgcacat gccttctcaa	4440
tgagactggc gacgagccct tccaatacaa aaactaatta gactttgagt gatcttgagc	4500
ctttcctagt tcatcccacc ccgccccaga gagatctttg tgaaggaacc ttacttctgt	4560
ggtgtgacat aattggacaa actacctaca gagatttaaa gctctaaggt aaatataaaa	4620


tttttaagtg	tataatgtgt	taaactactg	attctaattg	tttgtgtatt	ttagattcca	4680
acctatggaa						4740
tgctcagaag						4800
cctccaaaaa						4860
tttttgagtc						4920
aaggaaaaag						4980
agtaggcata						5040
gtgtctgcta	ttaataacta	tgctcaaaaa	ttgtgtacct	ttagcttttt	aatttgtaaa	5100
ggggttaata						5160
cacatttgta	gaggttttac	ttgctttaaa	aaacctccca	cacctccccc	tgaacctgaa	5220
acataaaatg	aatgcaattg	ttgttgttaa	cttgtttatt	gcagcttata	atggttacaa	5280
ataaagcaat						5340
tggtttgtcc						5400
gacggtgagg	cgcgcctact	gagtcattag	ggactttcca	atgggttttg	cccagtacat	5460
aaggtcaata	ggggtgaatc	aacaggaaag	tcccattgga	gccaagtaca	ctgagtcaat	5520
agggactttc	cattgggttt	tgcccagtac	aaaaggtcaa	tagggggtga	gtcaatgggt	5580
		acataaggtc				5640
caatttaatt	aaaacgccat	gtactttccc	accattgacg	tcaatgggct	attgaaacta	5700
atgcaacgtg	acctttaaac	ggtactttcc	catagctgat	taatgggaaa	gtaccgttct	5760
		ggaagtgaaa				5820
		tggcacgcat				5880
		cggtaccgtc				5940
		gtgagcaagc				6000
		ctggagggcg				6060
gctgcggcaa	gggcaacatc	ctgttcggca	accagctggt	gcagatccgc	gtgaccaagg	6120
gcgccccct	gcccttcgcc	ttcgacatcc	tgagccccgc	cttccagtac	ggcaaccgca	6180
		gacatcagcg				6240
		cgctacgagg				6300
tcaacctgat	cgaggagatg	ttcgtgtacc	gcgtggagta	caagggccgc	aacttcccca	6360
		aagaccatca				6420
		gtgggccagg				6480
		cgcaccctga				6540
ccgagtacca	cttcatccag	caccgcctgg	agaagaccta	cgtggaggac	ggcggcttcg	6600
tggagcagca	cgagaccgcc	atcgcccagc	tgaccagcct	gggcaagccc	ctgggcagcc	6660

• • •


1380

LBP30AND31.ST25.txt 6679 tgcacgagtg ggtgtaata <210> 2 <211> 8251 <212> DNA Hamster sp. <213> <220> misc_feature <221> Seq. ID. No. 3: circular plasmid GS vector p12.4 hCMVp-GFP /clone <223> <400> gaattcattg atcataatca gccataccac atttgtagag gttttacttg ctttaaaaaa 60 cctcccacac ctcccctga acctgaaaca taaaatgaat gcaattgttg ttgttaactt 120 gtttattgca gcttataatg gttacaaata aagcaatagc atcacaaatt tcacaaataa 180 agcatttttt tcactgcatt ctagttgtgg tttgtccaaa ctcatcaatg tatcttatca 240 tgtctggcgg ccgcgacctg caggcgcaga actggtaggt atggaagatc cctcgagatc 300 cattgtgctg gcggtaggcg agcagcgcct gcctgaagct gcgggcattc ccagtcagaa 360 atgagcgcca gtcgtcgtcg gctctcggca ccgaagtgct atgattctcc gccagcatgg 420 cttcggccag tgcgtcgagc agcgcccgct tgttcctgaa gtgccagtaa agcgccggct 480 gctgaacccc caaccgttcc gccagtttgc gtgtcgtcag accgtctacg ccgacctcgt 540 tcaacaggtc cagggcggca cggatcactg tattcggctg caactttgtc atgcttgaca 600 ctttatcact gataaacata atatgtccac caacttatca gtgataaaga atccgcgcca 660 gcacaatgga tctcgaggtc gagggatctc tagaggatcc atattcgcgg gcatcaccgg 720 cgccacaggt gcggttgctg gcgcctatat cgccgacatc accgatgggg aagatcgggc 780 tcgccacttc gggctcatga gcgcttgttt cggcgtgggt atggtggcag gccccgtggc 840 cgggggactg ttgggcgcca tctccttgca tgcaccattc cttgcggcgg cggtgctcaa 900 cggcctcaac ctactactgg gctgcttcct aatgcaggag tcgcataagg gagagcgtcg 960 acctcgggcc gcgttgctgg cgtttttcca taggctccgc ccccctgacg agcatcacaa 1020 aaatcgacgc tcaagtcaga ggtggcgaaa cccgacagga ctataaagat accaggcgtt 1080 tcccctgga agctcctcg tgcgctctcc tgttccgacc ctgccgctta ccggatacct 1140 gtccgccttt ctcccttcgg gaagcgtggc gctttctcat agctcacgct gtaggtatct 1200 cagttcggtg taggtcgttc gctccaagct gggctgtgtg cacgaacccc ccgttcagcc 1260 cgaccgctgc gccttatccg gtaactatcg tcttgagtcc aacccggtaa gacacgactt 1320

atcgccactg gcagcagcca ctggtaacag gattagcaga gcgaggtatg taggcggtgc



tacagagttc	ttgaagtggt	ggectaacta	cggctacact	agaagaacag	tatttggtat	1440
	ctgaagccag					1500
	gctggtagcg					1560
aaaaggatct	caagaagatc	ctttgatctt	ttctacgggg	tctgacgctc	agtggaacga	1620
aaactcacgt	taagggattt	tggtcatgag	attatcaaaa	aggatcttca	cctagatcct	1680
tttaaattaa	aaatgaagtt	ttaaatcaat	ctaaagtata	tatgagtaaa	cttggtctga	1740
cagttaccaa	tgcttaatca	gtgaggcacc	tatctcagcg	atctgtctat	ttcgttcatc	1800
catagttgcc	tgactccccg	tcgtgtagat	aactacgata	cgggagggct	taccatctgg	1860
ccccagtgct	gcaatgatac	cgcgagaccc	acgctcaccg	gctccagatt	tatcagcaat	1920
aaaccagcca	gccggaaggg	ccgagcgcag	aagtggtcct	gcaactttat	ccgcctccat	1980
ccagtctatt	aattgttgcc	gggaagctag	agtaagtagt	tcgccagtta	atagtttgcg	2040
caacgttgtt	gccattgcta	caggcatcgt	ggtgtcacgc	tcgtcgtttg	gtatggcttc	2100
attcagctcc	ggttcccaac	gatcaaggcg	agttacatga	tccccatgt	tgtgcaaaaa	2160
agcggttagc	tccttcggtc	ctccgatcgt	tgtcagaagt	aagttggccg	cagtgttatc	2220
actcatggtt	atggcagcac	tgcataattc	tcttactgtc	atgccatccg	taagatgctt	2280
ttctgtgact	ggtgagtact	caaccaagtc	attctgagaa	tagtgtatgc	ggcgaccgag	2340
ttgctcttgc	ccggcgtcaa	tacgggataa	taccgcgcca	catagcagaa	ctttaaaagt	2400
gctcatcatt	ggaaaacgtt	cttcggggcg	aaaactctca	aggatcttac	cgctgttgag	2460
atccagttcg	atgtaaccca	ctcgtgcacc	caactgatct	tcagcatctt	ttactttcac	2520
cagcgtttct	gggtgagcaa	aaacaggaag	gcaaaatgcc	gcaaaaaagg	gaataagggc	2580
gacacggaaa	tgttgaatac	tcatactctt	cctttttcaa	tattattgaa	gcatttatca	2640
gggttattgt	ctcatgagcg	gatacatatt	tgaatgtatt	tagaaaaata	aacaaatagg	2700
ggttccgcgc	acatttcccc	gaaaagtgcc	acctgacgtc	taagaaacca	ttattatcat	2760
gacattaacc	tataaaaata	ggcgtatcac	gaggccctga	tggctctttg	cggcacccat	2820
cgttcgtaat	gttccgtggc	accgaggaca	accctcaaga	gaaaatgtaa	tcacactggc	2880
tcaccttcgg	gtgggccttt	ctgcgtttat	aaggagacac	tttatgttta	agaaggttgg	2940
taaattcctt	gcggctttgg	cagccaagct	agatccggct	gtggaatgtg	tgtcagttag	3000
ggtgtggaaa	gtccccaggc	tccccagcag	gcagaagtat	gcaaagcatg	catctcaatt	3060
agtcagcaac	caggtgtgga	aagtccccag	gctccccagc	aggcagaagt	atgcaaagca	3120
	ttagtcagca					3180
ctccgcccag	ttccgcccat	tctccgcccc	atggctgact	aattttttt	atttatgcag	3240
aggccgaggo	cgcctcggc	: tctgagctat	tccagaagta	gtgaggaggc	ttttttggag	3300
gcctaggct1	ttgcaaaaag	ctagcttggg	gccaccgctc	agagcacctt	ccaccatggc	3360
cacctcagca	a agttcccact	tgaacaaaa	catcaagcaa	atgtacttgt	gcctgcccca	3420



		25.	. 50			
gggtgagaaa	gtccaagcca	tgtatatctg	ggttgatggt	actggagaag	gactgcgctg	3480
caaaacccgc	accctggact	gtgagcccaa	gtgtgtagaa	gagttacctg	agtggaattt	3540
tgatggctct	agtacctttc	agtctgaggg	ctccaacagt	gacatgtatc	tcagccctgt	3600
tgccatgttt	cgggacccct	tccgcagaga	tcccaacaag	ctggtgttct	gtgaagtttt	3660
caagtacaac	cggaagcctg	cagagaccaa	tttaaggcac	tcgtgtaaac	ggataatgga	3720
catggtgagc	aaccagcacc	cctggtttgg	aatggaacag	gagtatactc	tgatgggaac	3780
agatgggcac	ccttttggtt	ggccttccaa	tggctttcct	gggccccaag	gtccgtatta	3840
ctgtggtgtg	ggcgcagaca	aagcctatgg	cagggatatc	gtggaggctc	actaccgcgc	3900
ctgcttgtat	gctggggtca	agattacagg	aacaaatgct	gaggtcatgc	ctgcccagtg	3960
ggaactccaa	ataggaccct	gtgaaggaat	ccgcatggga	gatcatctct	gggtggcccg	4020
tttcatcttg	catcgagtat	gtgaagactt	tggggtaata	gcaacctttg	accccaagcc	4080
cattcctggg	aactggaatg	gtgcaggctg	ccataccaac	tttagcacca	aggccatgcg	4140
ggaggagaat	ggtctgaagc	acatcgagga	ggccatcgag	aaactaagca	agcggcaccg	4200
gtaccacatt	cgagcctacg	atcccaaggg	gggcctggac	aatgcccgtg	gtctgactgg	4260
gttccacgaa	acgtccaaca	tcaacgactt	ttctgctggt	gtcgccaatc	gcagtgccag	4320
catccgcatt	ccccggactg	tcggccagga	gaagaaaggt	tactttgaag	accgcggccc	4380
ctctgccaat	tgtgacccct	ttgcagtgac	agaagccatc	gtccgcacat	gccttctcaa	4440
tgagactggc	gacgagccct	tccaatacaa	aaactaatta	gactttgagt	gatcttgagc	4500
ctttcctagt	tcatcccacc	ccgccccaga	gagatctttg	tgaaggaacc	ttacttctgt	4560
ggtgtgacat	aattggacaa	actacctaca	gagatttaaa	gctctaaggt	aaatataaaa	4620
tttttaagtg	tataatgtgt	taaactactg	attctaattg	tttgtgtatt	ttagattcca	4680
acctatggaa	ctgatgaatg	ggagcagtgg	tggaatgcct	ttaatgagga	aaacctgttt	4740
tgctcagaag	aaatgccatc	tagtgatgat	gaggctactg	ctgactctca	acattctact	4800
cctccaaaaa	agaagagaaa	ggtagaagac	cccaaggact	ttccttcaga	attgctaagt	4860
tttttgagtc	atgctgtgtt	tagtaataga	actcttgctt	gctttgctat	ttacaccaca	4920
aaggaaaaag	ctgcactgct	atacaagaaa	attatggaaa	aatattctgt	aacctttata	4980
agtaggcata	acagttataa	tcataacata	ctgtttttc	ttactccaca	caggcataga	5040
gtgtctgcta	ttaataacta	tgctcaaaaa	ttgtgtacct	ttagcttttt	aatttgtaaa	5100
ggggttaata	aggaatattt	gatgtatagt	gccttgacta	gagatcataa	tcagccatac	5160
cacatttgta	gaggttttac	ttgctttaaa	aaacctccca	cacctccccc	tgaacctgaa	5220
acataaaatg	, aatgcaattg	, ttgttgttaa	cttgtttatt	gcagcttata	atggttacaa	5280
	agcatcacaa					5340
tggtttgtc	aaactcatca	atgtatctta	tcatgtctgg	atctagcttc	gtgtcaagga	5400
cggtgactg	agtgaataat	aaaatgtgtg	tttgtccgaa	atacgcgttt	tgagatttct	5460

PCT/EP2003/007946 WO 2004/009823 8/21

1. 12. 1. 15

gtcgccgact	aaattcatgt	cgcgcgatag	tggtgtttat	cgccgataga	gatggcgata	5520
ttggaaaaat	cgatatttga	aaatatggca	tattgaaaat	gtcgccgatg	tgagtttctg	5580
tgtaactgat	atcgccattt	ttccaaaagt	gatttttggg	catacgcgat	atctggcgat	5640
agcgcttata	tcgtttacgg	gggatggcga	tagacgactt	tggtgacttg	ggcgattctg	5700
tgtgtcgcaa	atatcgcagt	ttcgatatag	gtgacagacg	atatgaggct	atatcgccga	5760
tagaggcgac	atcaagctgg	cacatggcca	atgcatatcg	atctatacat	tgaatcaata	5820
ttggccatta	gccatattat	tcattggtta	tatagcataa	atcaatattg	gctattggcc	5880
attgcatacg	ttgtatccat	atcataatat	gtacatttat	attggctcat	gtccaacatt	5940
accgccatgt	tgacattgat	tattgactag	ttattaatag	taatcaatta	cggggtcatt	6000
agttcatagc	ccatatatgg	agttccgcgt	tacataactt	acggtaaatg	gcccgcctgg	6060
ctgaccgccc	aacgaccccc	gcccattgac	gtcaataatg	acgtatgttc	ccatagtaac	6120
gccaataggg	actttccatt	gacgtcaatg	ggtggagtat	ttacggtaaa	ctgcccactt	6180
ggcagtacat	caagtgtatc	atatgccaag	tacgccccct	attgacgtca	atgacggtaa	6240
atggcccgcc	tggcattatg	cccagtacat	gaccttatgg	gactttccta	cttggcagta	6300
catctacgta	ttagtcatcg	ctattaccat	ggtgatgcgg	ttttggcagt	acatcaatgg	6360
gcgtggatag	cggtttgact	cacggggatt	tccaagtctc	caccccattg	acgtcaatgg	6420
gagtttgttt	tggcaccaaa	atcaacggga	ctttccaaaa	tgtcgtaaca	actccgcccc	6480
attgacgcaa	atgggcggta	ggcgtgtacg	gtgggaggtc	tatataagca	gagctcgttt	6540
agtgaaccgt	cagatcgcct	ggagacgcca	tccacgctgt	tttgacctcc	atagaagaca	6600
ccgggaccga	tccagcctcc	gcggccggga	acggtgcatt	ggaacgcgga	ttccccgtgc	6660
caagagtgac	gtaagtaccg	cctatagagt	ctataggccc	acccccttgg	cttcttatgc	6720
atgctatact	gtttttggct	tggggtctat	acacccccgc	ttcctcatgt	tataggtgat	6780
ggtatagctt	agcctatagg	tgtgggttat	tgaccattat	tgaccactcc	cctattggtg	6840
acgatacttt	ccattactaa	tccataacat	ggctctttgc	cacaactctc	tttattggct	6900
atatgccaat	acactgtcct	tcagagactg	acacggactc	tgtattttta	caggatgggg	6960
tctcatttat	tatttacaaa	ttcacatata	caacaccacc	gtccccagtg	cccgcagttt	7020
ttattaaaca	taacgtggga	tctccacgcg	aatctcgggt	acgtgttccg	gacatgggct	7080
cttctccggt	agcggcggag	cttctacatc	cgagccctgc	tcccatgcct	ccagcgactc	7140
atggtcgctc	ggcagctcct	tgctcctaac	agtggaggcc	agacttaggc	acagcacgat	7200
gcccaccacc	accagtgtgc	cgcacaaggc	cgtggcggta	gggtatgtgt	ctgaaaatga	7260
gctcggggag	cgggcttgca	ccgctgacgc	atttggaaga	cttaaggcag	cggcagaaga	7320
agatgcaggc	agctgagttg	ttgtgttctg	ataagagtca	gaggtaactc	ccgttgcggt	7380
gctgttaacg	gtggagggca	gtgtagtctg	agcagtactc	gttgctgccg	cgcgcgccac	7440
cagacataat	agctgacaga	ctaacagact	gttcctttcc	atgggtcttt	tctgcagtca	7500

. 14

LBP30AND31.ST25.txt

ccgtccttga	cacgaagctt	gccgccacca	tggtgagcaa	gcagatcctg	aagaacaccg	7560
	gatcatgagc					7620
	gggctgcggc					7680
gcgtgaccaa	gggcgccccc	ctgcccttcg	ccttcgacat	cctgagcccc	gccttccagt	7740
acggcaaccg	caccttcacc	aagtaccccg	aggacatcag	cgacttcttc	atccagagct	7800
tccccgccgg	cttcgtgtac	gagcgcaccc	tgcgctacga	ggacggcggc	ctggtggaga	7860
tccgcagcga	catcaacctg	atcgaggaga	tgttcgtgta	ccgcgtggag	tacaagggcc	7920
gcaacttccc	caacgacggc	cccgtgatga	agaagaccat	caccggcctg	cagcccagct	7980
tcgaggtggt	gtacatgaac	gacggcgtgc	tggtgggcca	ggtgatcctg	gtgtaccgcc	8040
	caagttctac					8100
	ccccgagtac					8160
	cgtggagcag					8220
	cctgcacgag					8251

<210> 3

<211> 10369

<212> DNA

<213> Hamster sp.

<220>

<221> misc_feature

<223> seq. ID. No. 2: circular plasmid GS-minigene vector p 14.4 Deltam
 odulator (mod-) hCMVp-GFP /clone 6

<400> gaattcattg atcataatca gccataccac atttgtagag gttttacttg ctttaaaaaa 60 cctcccacac ctcccctga acctgaaaca taaaatgaat gcaattgttg ttgttaactt 120 gtttattgca gcttataatg gttacaaata aagcaatagc atcacaaatt tcacaaataa 180 agcatttttt tcactgcatt ctagttgtgg tttgtccaaa ctcatcaatg tatcttatca 240 tgtctggcgg ccgcgacctg caggcgcaga actggtaggt atggaagatc cctcgagatc 300 cattgtgctg gcggtaggcg agcagcgcct gcctgaagct gcgggcattc ccagtcagaa 360 atgagcgcca gtcgtcgtcg gctctcggca ccgaagtgct atgattctcc gccagcatgg 420 cttcggccag tgcgtcgagc agcgcccgct tgttcctgaa gtgccagtaa agcgccggct 480 gctgaacccc caaccgttcc gccagtttgc gtgtcgtcag accgtctacg ccgacctcgt 540 tcaacaggtc cagggcggca cggatcactg tattcggctg caactttgtc atgcttgaca 600

ctttatcact gataaacata	LB atatgtccac	P30AND31.ST caacttatca	25.txt gtgataaaga	atccgcgcca	660
gcacaatgga tctcgaggtc					720
cgccacaggt gcggttgctg					780
tcgccacttc gggctcatga					840
cgggggactg ttgggcgcca					900
cggcctcaac ctactactgg					960
acctcgggcc gcgttgctgg					1020
aaatcgacgc tcaagtcaga					1080
tcccctgga agctccctcg					1140
gtccgccttt ctcccttcgg					1200
cagttcggtg taggtcgttc					1260
cgaccgctgc gccttatccg					1320
atcgccactg gcagcagcca					1380
tacagagttc ttgaagtggt					1440
ctgcgctctg ctgaagccag					1500
acaaaccacc gctggtagcg					1560
aaaaggatct caagaagatc	ctttgatctt	ttctacgggg	tctgacgctc	agtggaacga	1620
aaactcacgt taagggattt					1680
tttaaattaa aaatgaagtt					1740
cagttaccaa tgcttaatca	gtgaggcacc	tatctcagcg	atctgtctat	ttcgttcatc	1800
catagttgcc tgactccccg	tcgtgtagat	aactacgata	cgggagggct	taccatctgg	1860
ccccagtgct gcaatgatac	cgcgagaccc	acgctcaccg	gctccagatt	tatcagcaat	1920
aaaccagcca gccggaaggg	ccgagcgcag	aagtggtcct	gcaactttat	ccgcctccat	1980
ccagtctatt aattgttgcc	gggaagctag	agtaagtagt	tcgccagtta	atagtttgcg	2040
caacgttgtt gccattgcta	caggcatcgt	ggtgtcacgo	tcgtcgtttg	gtatggcttc	2100
attcagctcc ggttcccaac	gatcaaggcg	agttacatga	tcccccatgt	tgtgcaaaaa	2160
agcggttagc tccttcggtc	ctccgatcgt	: tgtcagaagt	aagttggccg	cagtgttatc	2220
actcatggtt atggcagcad	tgcataatto	tcttactgtc	atgccatccg	taagatgctt	2280
ttctgtgact ggtgagtact	caaccaagto	: attctgagaa	tagtgtatgc	ggcgaccgag	2340
ttgctcttgc ccggcgtca	a tacgggataa	taccgcgcca	catagcagaa	ctttaaaagt	2400
gctcatcatt ggaaaacgt	t cttcggggc	g aaaactctca	a aggatcttac	cgctgttgag	2460
atccagttcg atgtaaccc	a ctcgtgcaco	caactgatct	tcagcatctt	ttactttcac	2520
cagcgtttct gggtgagca					2580
gacacggaaa tgttgaata	c tcatactct	t cctttttcaa	a tattattgaa	gcatttatca	2640

. 5

4.

4620

4680


LBP30AND31.ST25.txt gggttattgt ctcatgagcg gatacatatt tgaatgtatt tagaaaaata aacaaatagg 2700 ggttccgcgc acatttcccc gaaaagtgcc acctgacgtc taagaaacca ttattatcat 2760 gacattaacc tataaaaata ggcgtatcac gaggccctga tggctctttg cggcacccat 2820 cgttcgtaat gttccgtggc accgaggaca accctcaaga gaaaatgtaa tcacactggc 2880 tcaccttcgg gtgggccttt ctgcgtttat aaggagacac tttatgttta agaaggttgg 2940 taaattcctt gcggctttgg cagccaagct agatccagct ttttgcaaaa gcctaggcct 3000 ccaaaaaagc ctcctcacta cttctggaat agctcagagg ccgaggcggc ctcggcctct 3060 gcataaataa aaaaaattag tcagccatgg ggcggagaat gggcggaact gggcggagtt 3120 aggggcggga tgggcggagt taggggcggg actatggttg ctgactaatt gagatgcatg 3180 ctttgcatac ttctgcctgc tggggagcct ggggactttc cacacctggt tgctgactaa 3240 ttgagatgca tgctttgcat acttctgcct gctggggagc ctgggggactt tccacaccct 3300 aactgacaca cattccacag ggaagctagc ttggaattaa ttccccgccc ccttccaata 3360 caaaaactaa ttagactttg agtgatcttg agcctttcct agtttttgta ttggaagggc 3420 tcgtcgccag tctcattgag aaggcatgtg cggacgatgg cttctgtcac tgcaaagggg 3480 3540 aatgggagta gagccgactg cttgattccc acaccaatct cctcgccgct ctcacttcgc 3600 ctcgttctcg tggctcgtgg ccctgtccac cccgtccatc atcccgccgg ccaccgctca 3660 gagcaccttc caccatggcc acctcagcaa gttcccactt gaacaaaaac atcaagcaaa 3720 tgtacttgtg cctgccccag ggtgagaaag tccaagccat gtatatctgg gttgatggta 3780 ctggagaagg actgcgctgc aaaacccgca ccctggactg tgagcccaag tgtgtagaag 3840 agttacctga gtggaatttt gatggctcta gtacctttca gtctgagggc tccaacagtg 3900 acatgtatct cagccctgtt gccatgtttc gggacccctt ccgcagagat cccaacaagc 3960 tggtgttctg tgaagttttc aagtacaacc ggaagcctgc agagaccaat ttaaggcact 4020 cgtgtaaacg gataatggac atggtgagca accagcaccc ctggtttgga atggaacagg 4080 agtatactct gatgggaaca gatgggcacc cttttggttg gccttccaat ggctttcctg 4140 ggccccaagg tccgtattac tgtggtgtgg gcgcagacaa agcctatggc agggatatcg 4200 tggaggctca ctaccgcgcc tgcttgtatg ctggggtcaa gattacagga acaaatgctg 4260 aggtcatgcc tgcccagtgg gaactccaaa taggaccctg tgaaggaatc cgcatgggag 4320 atcatctctg ggtggcccgt ttcatcttgc atcgagtatg tgaagacttt ggggtaatag 4380 caacctttga ccccaagccc attcctggga actggaatgg tgcaggctgc cataccaact 4440 ttagcaccaa ggccatgcgg gaggagaatg gtctgaagta agtagctccc tctggaccat 4500 ctttattctc atggggtgga aggcctttgt gttagggttg ggaaaagttg gacttctcac 4560

aaactacatg ccatgctctt cgtgtttgtc ataagcctat cgttttgtac ccgttggaga

agtgacagta ctctaggaat agaattacag ctgtgatatg ggaaagttgt cacgtaggtt

(^ b

6720



gataactcct cattgtaaac agactgtaac tgccagagca gcgcttataa atcaacctaa



		1 R	P30AND31.ST	25.txt		
catttataag	atttcctctt	gacttgtttc	tttgtggttg	ggggaggaag	aaaaaaaaa	6780
gcgtgcagta	tttttttgtt	ccttcatttc	ctatcaaaag	aaaggggagt	ggttctgttt	6840
tgtttactcg	caaaataagc	tagcttatct	attggctttt	ctttttttt	ttttttttaa	6900
acgggctttt	tcttgtacct	ataatttggg	gtaaggtgtg	agagttttta	tagttttttg	6960
agacagggtc	ttggtgtata	cccttggctg	gcctggagct	aactatgtag	actgggctag	7020
cctttaactt	gcagttctgc	tttcaattag	ggtttataca	tttagtcttg	gcaattccta	7080
gttccacgtt	taatctcttt	acatttcaaa	gcagtgttat	ctgaagagtt	caggcgcaga	7140
gtcaattcaa	tagagttaca	caaaaaccta	aaaaacaagt	tttaaatacc	aagttatgtt	7200
ggcctggcca	cttttcacag	ctgtccacaa	ctcaatgtga	caaggctaca	aattggatat	7260
actagaattt	cctggtgatt	tggaacccct	gcttcatttc	ccggaaccag	ggcttttggt	7320
gacagtccta	gcttatcaga	ttatttaaaa	cagttactct	tcctgccctt	cttcctgaga	7380
cctttgtcca	gctgccatga	gccatctaca	cagtacttgc	ttccctgttg	aagtcactga	7440
aggcacatca	gcccaagaca	taaaggcttg	tcccggattc	actagcctgg	tgaacttgtg	7500
gttctctgat	gttttgtcct	gttttgttgt	gatttagtct	caaatttccc	agcctggttt	7560
gaaaatctgg	gctcccagcc	ttcaataagg	aggactacag	atatgtacga	ctgagccttg	7620
attccagcct	catgtttata	cgtctgtgct	cagctccctg	aaggttccag	tttgaaactc	7680
aataatccag	gggtcagaaa	gtcttgatct	tatccccaca	gtatggcacc	aagcctggct	7740
gagccttctg	acttagtctg	ccctgttgct	atttaagcac	ttttcttcac	taggctaaaa	7800
ataaaaggag	cttcctcctt	tgccatggcg	ctgtgcatga	taggaaaagg	tagctatcta	7860
ctagcatatt	aactccactg	tttttgcttt	gtgtgtttgg	tttttgagga	agggtctcaa	7920
ctgtgtatcc	ctggctggcc	tggccggatc	tagcttcgtg	tcaaggacgg	tgaggcgcgc	7980
caatattggc	tattggccat	tgcatacgtt	gtatccatat	cataatatgt	acatttatat	8040
tggctcatgt	ccaacattac	cgccatgttg	acattgatta	ttgactagtt	attaatagta	8100
atcaattacg	gggtcattag	ttcatagccc	atatatggag	ttccgcgtta	cataacttac	8160
ggtaaatggc	ccgcctggct	gaccgcccaa	cgacccccgc	ccattgacgt	caataatgac	8220
gtatgttccc	: atagtaacgc	caatagggac	tttccattga	cgtcaatggg	tggagtattt	8280
acggtaaact	gcccacttgg	cagtacatca	agtgtatcat	atgccaagta	cgcccctat	8340
tgacgtcaat	gacggtaaat	ggcccgcctg	gcattatgcc	cagtacatga	ccttatggga	8400
ctttcctact	tggcagtaca	tctacgtatt	agtcatcgct	attaccatgg	tgatgcggtt	8460
ttggcagtad	atcaatgggc	gtggatagcg	gtttgactca	cggggatttc	caagtctcca	8520
ccccattgad	gtcaatggga	gtttgttttg	gcaccaaaat	caacgggact	ttccaaaatg	8580
tcgtaacaad	tccgccccat	tgacgcaaat	gggcggtagg	cgtgtacggt	gggaggtcta	8640
tataagcaga	a gctcgtttag	tgaaccgtca	gatcgcctgg	agacgccatc	cacgctgttt	8700
tgacctcca	t agaagacacc	gggaccgatc	cagcctccgc	ggccgggaac	ggtgcattgg	8760

WO 2004/009823		FC1/E12	003/00/2
	14/21	•	
aacocooatt ccccotocca	LBP3OAND31.ST25.txt agagtgacgt aagtaccgcc tatagagtct	ataggcccac	8820
coccttoact tottatacat	gctatactgt ttttggcttg gggtctatac	accccgctt	8880
cctcatgtta taggtgatgg	tatagcttag cctataggtg tgggttattg	accattattg	8940
accacterer tattuutuac	gatactttcc attactaatc cataacatgg	ctctttgcca	9000
accaetetet tattggetat	atgccaatac actgtccttc agagactgac	acggactctg	9060
	tcatttatta tttacaaatt cacatataca		9120

tatttttaca ggatggggtc tcatttatta tttacaaatt cacat ccccagtgcc cgcagttttt attaaacata acgtgggatc tccacgcgaa tctcgggtac 9180 gtgttccgga catgggctct tctccggtag cggcggagct tctacatccg agccctgctc 9240 ccatgcctcc agcgactcat ggtcgctcgg cagctccttg ctcctaacag tggaggccag 9300 acttaggcac agcacgatgc ccaccaccac cagtgtgccg cacaaggccg tggcggtagg 9360 gtatgtgtct gaaaatgagc tcggggagcg ggcttgcacc gctgacgcat ttggaagact 9420 taaggcagcg gcagaagaag atgcaggcag ctgagttgtt gtgttctgat aagagtcaga 9480 ggtaactccc gttgcggtgc tgttaacggt ggagggcagt gtagtctgag cagtactcgt 9540 tgctgccgcg cgcgccacca gacataatag ctgacagact aacagactgt tcctttccat 9600 gggtcttttc tgcagtcacc gtccttgaca cgaagcttgc cgccaccatg gtgagcaagc 9660 agatcctgaa gaacaccggc ctgcaggaga tcatgagctt caaggtgaac ctggagggcg 9720 tggtgaacaa ccacgtgttc accatggagg gctgcggcaa gggcaacatc ctgttcggca 9780 accagctggt gcagatccgc gtgaccaagg gcgccccctt gcccttcgcc ttcgacatcc 9840 tgagccccgc cttccagtac ggcaaccgca ccttcaccaa gtaccccgag gacatcagcg 9900 acttcttcat ccagagcttc cccgccggct tcgtgtacga gcgcaccctg cgctacgagg 9960 acggcggcct ggtggagatc cgcagcgaca tcaacctgat cgaggagatg ttcgtgtacc 10020 gcgtggagta caagggccgc aacttcccca acgacggccc cgtgatgaag aagaccatca 10080 ccggcctgca gcccagcttc gaggtggtgt acatgaacga cggcgtgctg gtgggccagg 10140 tgatcctggt gtaccgcctg aacagcggca agttctacag ctgccacatg cgcaccctga 10200 tgaagagcaa gggcgtggtg aaggacttcc ccgagtacca cttcatccag caccgcctgg 10260 agaagaccta cgtggaggac ggcggcttcg tggagcagca cgagaccgcc atcgcccagc 10320

tgaccagcct gggcaagccc ctgggcagcc tgcacgagtg ggtgtaata

<210> 4

٠١٠

<211> 13535

<212> DNA

<213> Hamster sp.

<220>

<221> misc_feature

e b



LBP3OAND31.ST25.txt


<223> Seq. ID. No. 1: circular plasmid GS + IgG 2A hot spot targetting vector pEE 15.1 hCMVp-GFP /clone 11

<400> 4 gaattcattg	atcataatca	gccataccac	atttgtagag	gttttacttg	ctttaaaaaa	60
cctcccacac	ctccccctga	acctgaaaca	taaaatgaat	gcaattgttg	ttgttaactt	120
gtttattgca	gcttataatg	gttacaaata	aagcaatagc	atcacaaatt	tcacaaataa	180
agcattttt	tcactgcatt	ctagttgtgg	tttgtccaaa	ctcatcaatg	tatcttatca	240
tgtctggcgg	ccgcgacctg	caggcgcaga	actggtaggt	atggaagatc	cctcgagatc	300
cattgtgctg	gcggtaggcg	agcagcgcct	gcctgaagct	gcgggcattc	ccagtcagaa	360
atgagcgcca	gtcgtcgtcg	gctctcggca	ccgaagtgct	atgattctcc	gccagcatgg	420
cttcggccag	tgcgtcgagc	agcgcccgct	tgttcctgaa	gtgccagtaa	agcgccggct	480
gctgaacccc	caaccgttcc	gccagtttgc	gtgtcgtcag	accgtctacg	ccgacctcgt	540
tcaacaggtc	tagggcggca	cggatcactg	tattcggctg	caactttgtc	atgcttgaca	600
ctttatcact	gataaacata	atatgtccac	caacttatca	gtgataaaga	atccgcgcca	660
gcacaatgga	tctcgaggtc	gagggatctc	tagaggatcc	atattcgcga	atatgccggc	720
atcaccggcg	ccacaggtgc	ggttgctggc	gcctatatcg	ccgacatcac	cgatggggaa	780
gatcgggctc	gccacttcgg	gctcatgagc	gcttgtttcg	gcgtgggtat	ggtggcaggc	840
ccgtggccgg	gggactgttg	ggcgccatct	ccttgcatgc	accattcctt	gcggcġgcgg	900
tgctcaacgg	cctcaaccta	ctactgggct	gcttcctaat	gcaggagtcg	cataagggag	960
agcgtcgagt	cctccgtgtt	cgaagcgatc	cctgtccagt	ggtgtgcaca	ccttcccagc	1020
tgtcctgcag	tctgacctct	acaccctcag	cagctcagtg	actgtaacct	cgagcacctg	1080
gcccagccag	tccatcacct	gcaatgtggc	ccacccggca	agcagcacca	aggtggacaa	1140
gaaaattggt	gaggaaaaca	aggggagtag	aggttcacaa	gtgattagtc	taaggcctta	1200
gcctagctag	accagccagg	atcagcagcc	atcaccaaaa	atgggaactt	ggcccagaag	1260
agaaggagat	actgactgtg	actccctctt	ggaaacttct	aactatgacc	acctaccttc	1320
aaggtcatga	tcctctagga	tagatgtcct	tggtcatttc	caggatcatc	ctgacctaag	1380
gccataccca	gggacaaagt	ccctggtttg	gtgccttttc	tccttcaaac	ttgagtaacc	1440
cccagccttc	tctctgcaga	gcccagaggg	cccacaatca	agccctgtcc	tccatgcaaa	1500
tgcccaggta	agtcactaga	ccagagctcc	acccgggaga	atggtaagtg	ctgtaaacat	1560
ccctgcacta	gaggataagc	catgtacaga	tccatttcca	tctctcctca	tcagcaccta	1620
acctcttggg	tggaccatcc	gtcttcatct	tccctccaaa	gatcaaggat	gtactcatga	1680
tctccctgag	ccccatagtc	acatgtgtgg	tggtggatgt	gagcgaggat	gacccagatg	1740
tccagatcag	ctggtttgtg	aacaacgtgg	aagtacacac	agctcagaca	caaacccata	1800
gagaggatta	caacagtact	ctccgggtgg	tcagtgccct	ccccatccag	caccaggact	1860

6.36 - 31




ggatgagtgg						1920
		aaaggtgaga				1980
gggcataagg	ataaaggtct	gtgtggacag	ccttctgctt	cagccatgac	ctttgtgtat	2040
gtttctaccc	tcacagggtc	agtaagagct	ccacaggtat	atgtcttgcc	tccaccagaa	2100
gaagagatga	ctaagaaaca	ggtcactctg	acctgcatgg	tcacagactt	catgcctgaa	2160
gacatttacg	tggagtggac	caacaacggg	aaaacagagc	taaactacaa	gaacactgaa	2220
ccagtcctgg	actctgatgg	ttcttacttc	atgtacagca	agctgagagt	ggaaaagaag	2280
aactgggtgg	aaagaaatag	ctactcctgt	tcagtggtcc	acgagggtct	gcacaatcac	2340
cacacgacta	agagcttctc	ccggactccg	ggtaaatgag	ctcagcaccc	acaaaactct	2400
caggtccaaa	gagacaccca	cactcatctc	catgcttccc	ttgtataaat	aaagcaccca	2460
ccaatgcctg	ggaccatgta	aaactgtcct	ggttctttcc	aaggtataga	gcatagctca	2520
caggctgata	tttctggcca	gggttggagg	acagccttgt	ctataggaag	agaatgaggt	2580
ttttgcactg	caggactcag	agctcattag	ttatcctgcc	ttggagtgtt	ggggcttggc	2640
tttaggcagt	gccttttcct	tgccttccta	cgaaccagca	gctgccatac	atagagataa	2700
tcctaggaag	cctcaaatgg	agaaggacac	aaacccacct	ccctcaggct	gttcctctat	2760
cccggcccca	cttctttacc	taggggtttc	tctgagtcta	ttgtggagtt	acacatggcc	2820
aggggcattc	cagagaccct	tgtcatccat	acactcaact	caggcagctt	tgcacaaaca	2880
aagtctgcac	acccatacag	atggctcact	cttgcctgtg	ccatgtaggg	ctgaggcaca	2940
tggctcttgc	tgccccaagg	gagggactat	tagatagcca	cactcatgct	gaatcctggc	3000
ccattcaaat	tagcctgctg	aacaccatcc	agtccatata	gcacatgtat	ccacatgcac	3060
gtgtgcacaa	aacgcattta	atacactggg	acaacaattc	tgtgccctgc	acagcaccta	3120
tatccagcaa	tgtatcacca	tacacacgac	caaaaaaatt	caatgcccac	gtttctgcca	3180
tcacaaacag	acacatcttt	cctctctgtg	gccactgcat	tatatgctca	acacaagacc	3240
tctgaagcca	gatccatctc	tggcacctcg	gggtcatgct	tcaaccccac	atgaattatg	3300
caaaccatag	ccataatggt	ctgaatcact	tcacactggg	atgttcccaa	gttcaggcaa	3360
gacgagccac	aggctctgct	gatgactgaa	ggacagcaaa	gggtcagtcc	agctgtatag	3420
ccactgttga	cctgggtcac	aggccctgct	gaccctccac	cttctcctgt	actgaaggaa	3480
tgaaagatga	gacaagcata	gagggcactt	gaataatcca	ggtcactctg	aggtccaccc	3540
aaggcattat	tggactcagg	tgggaagctg	agactggtgt	cccagaggga	aaggaaggaa	3600
agcaggcccc	ggggagggtc	tgctgtccca	gtcaggctgg	agatctctcc	tctgaatcca	3660
tgcagacatg	tctgcctcac	agggaatctc	tcccagcacc	aaccatgttg	ggacaaacac	3720
tgactgtcct	ctctgttcag	ggctagacct	ggatgatgtc	tgtgctgagg	cccaggacgg	3780
ggagctggac	ggcctctgga	cgaccatcac	catcttcatc	agcctcttcc	tgctcagcgt	3840
gtgctacagc	gcctctgtca	cactcttcaa	ggttggcact	gtctcccacc	ctctgctgtg	3900



atggctacac	tgaccacaaa	atgtcctctc	actcctcccc	agatgtagta	ggacgttact	3960
ttgctgcccc	tactctgtcc	cacacaccat	ttcctccatt	ccctgagcca	tcccacattg	4020
ttctatgtga	ctccacattg	tgtcccatac	agtctgccct	tctgtctctc	tggctgtcct	4080
gcgtgatcct	gatactgtct	tatgagacca	aacctccttg	cattccacac	tagccttcat	4140
gaggttcaat	gctgtcttac	acacaatccc	ctcagcctca	ccatggctca	aggtactctg	4200
tgagctatcc	tcataccatc	tccacctcaa	ctcccacaat	atctccactc	tgacccctcc	4260
catacccagt	ctcctacctg	tatgaaggga	attgaaggag	agacaggtcg	acctctgtct	4320
ttcccacaga	ttggagggtc	tgagcatggg	cgtggtctct	gactttctct	cacttcccca	4380
caggtaaagt	ggatcttctc	ctctgtggtg	gagctgaagc	agacgatctc	ccctgactac	4440
agaaacatga	ttgggcaggg	agcctaggcc	acttcctctg	ggatcagaag	agcttcctag	4500
gccctgcaga	agcccatcca	tcctactgtg	cagcctaaca	gggaggccac	actctagccc	4560
tatgactctc	tgatcagaac	tcccatggtc	tcctctttgg	aggaccacgt	gcagtgcagg	4620
ctttgcccag	acctaaacac	ttccacagca	gtcgccagat	atctaactac	tccggaccag	4680
aagaaccatc	tccttccaaa	ccagcactag	ggatctgaga	tctcagaatg	tttgcctaag	4740
aagagctgga	aatccaggct	tcctgtgttc	tgctacaagg	acatcagcct	ggatttgacc	4800
tggaccacac	attttcatct	aaatgagttt	tccacaaagg	acacgtttca	gatccttgaa	4860
tgagacctct	acatggaaga	ccagagtcac	tatacccaaa	ggtcactctg	tatccttgca	4920
ccagctatac	tggacagctt	ccttcctggt	acttcagtga	ccctggctga	ggaaaggatc	4980
tgtgacctca	actgtttgga	gagcctctgg	aagatgtagt	cttctcttcc	tgctaccacc	5040
aacatgctgg	atctcagatg	cagaatccaa	tccacagaca	ccactgacca	cacaacctga	5100
agacaaggcc	attgccacct	ccacagagat	gccatccaca	ctctgtggag	aaataaggag	5160
tgctttgtgc	agcctctgca	aagctctggc	agggattaga	gtatacacac	tgagtactga	-5220
ctaggtgacc	aggcagaaaa	acctccagga	gaaggaacaa	tgggggagag	atgtgaacag	5280
atagttagaa	aaagcatggt	gtcacaggtc	tgctctgtgg	actgatttcc	agattggacc	5340
acctacagca	gaaaccatcg	gttgcagtgg	caatctagga	ggaccaacct	ggaataggag	5400
ggctgctgtg	gtcaatggag	agtagacctg	tatctatttc	tccactgcct	cttatgacca	5460
ataagaagco	agagtctcca	gacagaaaga	aagaaagaaa	gaaagaaaga	aagaaagaaa	5520
gaaagagaga	gagagagaga	gagagagaga	gaggaaggaa	ggaaggaagg	aaggaaggaa	5580
ggaaggaagg	aggaggagga	ggaggaggag	gaggaggaga	gagagagaga	gagagaga	5640
gagagagaga	gagcaccagc	ttttctgtga	ctggaaggaa	atgcttagag	agcttggatc	5700
tttaaagctt	cttttttcta	gagaccatga	atgtctttgt	tctctctctc	tctctctctc	5760
tctctctct	tctctctctc	tctctctgtg	tgtgtgtgtg	tgtgtgtgtg	tgtgtgtgcg	5820
tgcatgcacg	, ctattgttt	ggcatttgaa	acaataaaac	attcttttaa	tattctgtat	5880
ctcatggtto	cccttctgtg	tggatcagcc	ctaacaccca	ggaacagggg	acaataaaca	5940


L34 1 3





gaccacagcc	atgtacagcc	ttctacctcc	cttctggttc	tgacctccca	gaggtccctc	6000
agtgggcccc	tcacagctgg	gtttcttccc	tggcagtgcc	accaagagct	caggcacctc	6060
tgagctggag	gctgtcctga	tgccataggc	aggctatgga	gcagagatga	tgaccacggt	6120
ggactccagg	tgagccaggc	aaagcctccc	atgccagaag	agaagcgtgt	ggtactcact	6180
ggcctcgggc	tgctacggat	tcagcaaaga	gcatggatcg	cttcgaagcc	tccaagctcg	6240
acctcgggcc	gcgttgctgg	cgtttttcca	taggctccgc	cccctgacg	agcatcacaa	6300
aaatcgacgc	tcaagtcaga	ggtggcgaaa	cccgacagga	ctataaagat	accaggcgtt	6360
tcccctgga	agctccctcg	tgcgctctcc	tgttccgacc	ctgccgctta	ccggatacct	6420
gtccgccttt	ctcccttcgg	gaagcgtggc	gctttctcaa	tgctcacgct	gtaggtatct	6480
cagttcggtg	taggtcgttc	gctccaagct	gggctgtgtg	cacgaacccc	ccgttcagcc	6540
cgaccgctgc	gccttatccg	gtaactatcg	tcttgagtcc	aacccggtaa	gacacgactt	6600
atcgccactg	gcagcagcca	ctggtaacag	gattagcaga	gcgaggtatg	taggcggtgc	6660
tacagagttc	ttgaagtggt	ggcctaacta	cggctacact	agaaggacag	tatttggtat	6720
ctgcgctctg	ctgaagccag	ttaccttcgg	aaaaagagtt	ggtagctctt	gatccggcaa	6780
acaaaccacc	gctggtagcg	gtggttttt	tgtttgcaag	cagcagatta	cgcgcagaaa	6840
aaaaggatct	caagaagatc	ctttgatctt	ttctacgggg	tctgacgctc	agtggaacga	6900
aaactcacgt	taagggattt	tggtcatgag	attatcaaaa	aggatcttca	cctagatcct	6960
tttaaattaa	aaatgaagtt	ttaaatcaat	ctaaagtata	tatgagtaaa	cttggtctga	7020
cagttaccaa	tgcttaatca	gtgaggcacc	tatctcagcg	atctgtctat	ttcgttcatc	7080
catagttgcc	tgactccccg	tcgtgtagat	aactacgata	cgggagggct	taccatctgg	7140
ccccagtgct	gcaatgatac	cgcgagaccc	acgctcaccg	gctccagatt	tatcagcaat	7200
	gccggaaggg					7260
	aattgttgcc					7320
caacgttgtt	gccattgcta	caggcatcgt	ggtgtcacgc	tcgtcgtttg	gtatggcttc	7380
	ggttcccaac					7440
	tccttcggtc					7500
	atggcagcac					7560
	ggtgagtact					7620
	ccggcgtcaa					7680
	ggaaaacgtt					7740
	atgtaaccca					7800
	gggtgagcaa					7860
	tgttgaatac					7920
gggttattgt	: ctcatgagcg	gatacatatt	tgaatgtatt	tagaaaaata	aacaaatagg	7980

₩O 2004/009823



LBP30AND31.ST25.txt

19/21

ggttccgcgc acatttcccc	gaaaagtgcc	acctgacgtc	taagaaacca	ttattatcat	8040
gacattaacc tataaaaata					8100
cgttcgtaat gttccgtggc					8160
tcaccttcgg gtgggccttt					8220
taaattcctt gcggctttgg					8280
agggtgtgga aagtccccag					8340
ttagtcagca accaggtgtg					8400
catgcatctc aattagtcag	caaccatagt	cccgccccta	actccgccca	tcccgcccct	8460
aactccgccc agttccgccc	attctccgcc	ccatggctga	ctaatttttt	ttatttatgc	8520
agaggccgag gccgcctcgg					8580
aggcctaggc ttttgcaaaa	agctagcttg	gggccaccgc	tcagagcacc	ttccaccatg	8640
gccacctcag caagttccca	cttgaacaaa	aacatcaagc	aaatgtactt	gtgcctgccc	8700
cagggtgaga aagtccaagc	catgtatatc	tgggttgatg	gtactggaga	aggactgcgc	8760
tgcaaaaccc gcaccctgga	ctgtgagccc	aagtgtgtag	aagagttacc	tgagtggaat	8820
tttgatggct ctagtacctt	tcagtctgag	ggctccaaca	gtgacatgta	tctcagccct	8880
gttgccatgt ttcgggaccc	cttccgcaga	gatcccaaca	agctggtgtt	ctgtgaagtt	8940
ttcaagtaca accggaagcc	tgcagagacc	aatttaaggc	actcgtgtaa	acggataatg	9000
gacatggtga gcaaccagca					9060
acagatgggc acccttttgg	ttggccttcc	aatggctttc	ctgggcccca	aggtccgtat	9120
tactgtggtg tgggcgcaga	caaagcctat	ggcagggata	tcgtggaggc	tcactaccgc	9180
gcctgcttgt atgctggggt			•		9240
tgggaactcc aaataggacc	ctgtgaagga	atccgcatgg	gagatcatct	ctgggtggcc	9300
cgtttcatct tgcatcgagt	atgtgaagac	tttggggtaa	tagcaacctt	tgaccccaag	9360
cccattcctg ggaactggaa	tggtgcaggc	tgccatacca	actttagcac	caaggccatg	9420
cgggaggaga atggtctgaa	gcacatcgag	gaggccatcg	agaaactaag	caagcggcac	9480
cggtaccaca ttcgagccta	cgatcccaag	gggggcctgg	acaatgcccg	tggtctgact	9540
gggttccacg aaacgtccaa	catcaacgac	ttttctgctg	gtgtcgccaa	tcgcagtgcc	9600
agcatccgca ttccccggac	tgtcggccag	gagaagaaag	gttactttga	agaccgcggc	9660
ccctctgcca attgtgaccc	ctttgcagtg	acagaagcca	tcgtccgcac	atgccttctc	9720
aatgagactg gcgacgagcc	cttccaatac	aaaaactaat	tagactttga	gtgatcttga	9780
gcctttccta gttcatccca	cccgcccca	gagagatctt	tgtgaaggaa	ccttacttct	9840
gtggtgtgac ataattggac	aaactaccta	cagagattta	aagctctaag	gtaaatataa	9900
aatttttaag tgtataatgt	gttaaactac	: tgattctaat	tgtttgtgta	ttttagattc	9960
caacctatgg aactgatgaa	tgggagcagt	ggtggaatgc	ctttaatgag	gaaaacctgt	10020



LBP30AND31.ST25.txt

20/21

tttgctcag	ja a	igaaatgcca	tctagtgatg	atgaggctac	tgctgactct	caacattcta	10080
ctcctccaa	ia a	aagaagaga	aaggtagaag	accccaagga	ctttccttca	gaattgctaa	10140
atttttqa	la 1	tcatqctqtg	tttagtaata	gaactcttgc	ttgctttgct	atttacacca	10200
caaaggaaa	aa a	agctgcactg	ctatacaaga	aaattatgga	aaaatattct	gtaaccttta	10260
taaqtaqq	:a 1	taacagttat	aatcataaca	tactgttttt	tcttactcca	cacaggcata	10320
			tatgctcaaa				10380
			ttgatgtata				10440
			acttgcttta				10500
			tgttgttgtt				10560
			aaatttcaca				10620
			caatgtatct				10680
			taataaaatg				10740
			atgtcgcgcg				10800
			ttgaaaatat				10860
			atttttccaa				10920
			acgggggatg				10980
			cagtttcgat				11040
			ctggcacatg				11100
			ttattcattg				11160
ggccattg	ca	tacgttgtat	ccatatcata	atatgtacat	ttatattggc	tcatgtccaa	11220
			tgattattga				11280
cattagtt	ca	tagcccatat	atggagttcc	gcgttacata	acttacggta	aatggcccgc	11340
ctggctga	cc	gcccaacgac	ccccgcccat	tgacgtcaat	aatgacgtat	gttcccatag	11400
taacgcca	at	agggactttc	cattgacgtc	aatgggtgga	gtatttacgg	taaactgccc	11460
						gtcaatgacg	11520
						cctacttggc	11580
						cagtacatca	11640
						attgacgtca	11700
						aacaactccg	11760
						agcagagctc	11820
						ctccatagaa	11880
						cggattcccc	11940
						ttggcttctt	12000
atgcatg	cta	tactgtttt	t ggcttggggt	ctatacacco	ccgcttcct	atgttatagg	12060



tgatggtata	gcttagccta	taggtgtggg	ttattgacca	ttattgacca	ctcccctatt	12120
gotgacgata	ctttccatta	ctaatccata	acatggctct	ttgccacaac	tctctttatt	12180
ggctatatgc	caatacactg	tccttcagag	actgacacgg	actctgtatt	tttacaggat	12240
agaatctcat	ttattattta	caaattcaca	tatacaacac	caccgtcccc	agtgcccgca	12300
ottttatta	aacataacgt	gggatctcca	cgcgaatctc	gggtacgtgt	tccggacatg	12360
gactcttctc	cggtagcggc	ggagcttcta	catccgagcc	ctgctcccat	gcctccagcg	12420
actcatqqtc	gctcggcagc	tccttgctcc	taacagtgga	ggccagactt	aggcacagca	12480
cgatgcccac	caccaccagt	gtgccgcaca	aggccgtggc	ggtagggtat	gtgtctgaaa	12540
atgagctcgg	ggagcgggct	tgcaccgctg	acgcatttgg	aagacttaag	gcagcggcag	12600
aagaagatgc	aggcagctga	gttgttgtgt	tctgataaga	gtcagaggta	actcccgttg	12660
caatactatt	aacggtggag	ggcagtgtag	tctgagcagt	actcgttgct	gccgcgcgcg	12720
ccaccagaca	taatagctga	cagactaaca	gactgttcct	ttccatgggt	cttttctgca	12780
atcaccatca	ttgacacgaa	gcttgccgcc	accatggtga	gcaagcagat	cctgaagaac	12840
acconctoc	aggagatcat	gagcttcaag	gtgaacctgg	agggcgtggt	gaacaaccac	12900
atattcacca	tggagggctg	cggcaagggc	aacatcctgt	tcggcaacca	gctggtgcag	12960
atccgcgtga	ccaagggcgc	cccctgccc	ttcgccttcg	acatcctgag	cccgccttc	13020
cagtacggca	accgcacctt	caccaagtac	cccgaggaca	tcagcgactt	cttcatccag	13080
agcttcccc	ccggcttcgt	gtacgagcgc	accctgcgct	acgaggacgg	cggcctggtg	13140
gagatccgca	gcgacatcaa	cctgatcgag	gagatgttcg	tgtaccgcgt	ggagtacaag	13200
aaccacaact	tccccaacga	cggccccgtg	atgaagaaga	ccatcaccgg	cctgcagccc	13260
agettegag	tggtgtacat	gaacgacggc	gtgctggtgg	gccaggtgat	cctggtgtac	13320
cocctoaaca	gcggcaagtt	ctacagctgc	: cacatgcgca	ccctgatgaa	gagcaagggc	13380
gtagtgaag	g acttccccga	gtaccactto	atccagcacc	gcctggagaa	gacctacgtg	13440
gaggacggc	g gcttcgtgga	gcagcacgag	accgccatcg	cccagctgac	cagcctgggc	13500
	g gcagcctgca					13535
		_				